


Do oil-producing crops need extra manganese or just enough?

Manganese is a workhorse micronutrient in plants. It is central to photosystem II, essential for the water splitting chemistry, and a cofactor for several enzymes. Given its importance, plants that produce energetically expensive compounds – like oils – might require more of it to run their machinery, so the threshold question is simple: do oilseed or essential oil crops require manganese above what non oil-producers need, or do they just need standard sufficiency with no premium for “oil production status”?



A manganese sulfate crystal. One of the most commonly used salts to supplement Mn in agriculture.

What the literature actually supports

Recent reviews agree on fundamentals. Plant Mn requirements are driven by core physiology like photosynthesis and redox

balance, not by whether a crop partitions carbon to oil, starch or protein. There is no general evidence for a higher Mn setpoint in oil-producing species as a class. Instead, yield and quality respond to correcting deficiency and avoiding toxicity, the same rule that governs non oil-producing crops [\(1\)](#), [\(2\)](#).

Oilseeds

- **Soybean.** Classic work shows severe Mn deficiency reduces seed oil percentage. Once deficiency is corrected, pushing Mn higher does not increase oil; excess Mn depresses growth and yield. In other words, soybean needs adequate Mn, not extra because it is an oilseed [\(3\)](#), [\(4\)](#).
- **Canola/rapeseed.** Liming-induced Mn deficiency is common on high pH soils. Foliar Mn corrects deficiency and restores yield, but applications on adequate plants do not increase oil or seed yield. Again, the benefit is deficiency correction, not a special oil-crop premium [\(5\)](#).

Essential oil crops

- **Water mint (*Mentha aquatica*).** In solution culture, applying 100 μ M Mn sulfate, which is ~5.5 ppm Mn, increased leaf glandular trichome density and essential oil yield relative to a lower Mn background. This shows Mn can modulate secondary metabolism when the baseline is low, but it does not prove that mint requires Mn above typical sufficiency ranges; it shows that deficiency or marginal supply limits oil yield and composition [\(6\)](#).
- **Feverfew (*Tanacetum parthenium*).** Varying Mg and Mn in controlled media shifted essential oil profiles. Mn

interacted with Mg to alter monoterpene vs sesquiterpene proportions, again indicating composition sensitivity under limited or imbalanced supply rather than a universal need for “extra Mn” [\(7\)](#).

Soilless and hydroponic angle

Hydroponics removes soil redox chemistry, so Mn availability is governed by solution concentration, chelation and pH. Reviews emphasize that plants still follow the same homeostatic rules; oil status does not change the Mn target. In recirculating systems, Mn can drift due to adsorption, precipitation at higher pH and plant uptake, which explains sporadic deficiency in otherwise balanced recipes. Correct the drift and the symptoms resolve; adding more than sufficiency is unnecessary and risks toxicity, especially at low pH [\(1\)](#), [\(2\)](#).

Evidence summary

Crop	System	Mn supplementation rate (ppm Mn)	Outcome on oil yield or composition	Take-home	Study
Soybean	Sand/solution culture	Not specified here in ppm	Severe Mn deficiency lowered seed oil; correcting deficiency restored yield but extra Mn gave no benefit	Adequacy matters, excess does not help	(3) , (4)

Crop	System	Mn supplementation rate (ppm Mn)	Outcome on oil yield or composition	Take-home	Study
Canola	Field, calcareous soils	Foliar Mn, rate study	Yield gains only where tissue was Mn-deficient; no gain in Mn-sufficient stands	Target deficiency, not blanket “oil-crop” boosts	(5)
Water mint	Nutrient solution	~5.5	Increased trichome density and essential oil yield from a low-Mn baseline	Adequate Mn is required for E0 biosynthesis; no proof of supra-sufficiency need	(6)
Feverfew	Controlled media	Varied Mn, ppm not reported	Mn with Mg shifted monoterpene vs sesquiterpene proportions	Composition responds to Mn status; optimize for sufficiency	(7)

Tissue composition: are oil plants different?

Authoritative reviews catalog Mn uptake, transport and intracellular allocation across species. None propose distinct Mn sufficiency thresholds based solely on oil production. The drivers are photosynthetic demand, transporter regulation and rhizosphere chemistry. Oilseed and essential oil crops display the same deficiency symptoms and toxicity risks as other species. Practically, tissue targets should be set by species-specific sufficiency ranges and growth stage, not by “oil producer” status [\(1\)](#), [\(2\)](#).

Practical stance for soilless growers

1. **Aim for sufficiency, verify with tissue tests.** If chlorosis and interveinal speckling suggest Mn deficiency and tissue Mn is low, bring solution Mn up to a normal range and adjust pH. Do not chase extra Mn for oil content once sufficiency is confirmed [\(5\)](#).
2. **Watch pH and redox.** Slight pH rises or oxidizing conditions can drop available Mn even when total Mn dosing looks fine. Correct pH and renew chelates before increasing Mn concentration [\(1\)](#).
3. **Expect composition shifts near the margins.** In mint and feverfew, Mn status influenced essential oil profile when supply was marginal. That is a signal to maintain adequacy, not a license to overapply [\(6\)](#), [\(7\)](#).

Bottom line

There is no broad academic support for supplementing manganese above normal sufficiency just because a crop produces oil. The consistent finding is boring but useful: correct Mn deficiency and keep supply in a normal, pH-stable window. Oilseed yield and essential oil profiles suffer when Mn is low, and they recover when Mn is adequate. Beyond that, extra Mn does not buy more oil and can cost you growth.