


Ascorbic Acid as a Biostimulant: Enhancing Yield and Quality in Hydroponic Systems

The search for sustainable biostimulants to enhance crop productivity has led researchers to investigate ascorbic acid (vitamin C) as a promising alternative to synthetic growth regulators. This natural antioxidant compound has shown remarkable potential in improving both yield and quality parameters in hydroponic and soilless cultivation systems.



Vitamin C
 $C_6H_8O_6$



Model representation of Ascorbic Acid (Vitamin C)

Understanding Ascorbic Acid as a Biostimulant

Ascorbic acid functions as a [\(1\)](#) multifunctional non-enzymatic antioxidant that plays crucial roles in plant physiology beyond its traditional vitamin C function. In hydroponic systems, ascorbic acid applications can modulate several key physiological processes including photosynthesis, antioxidant defense mechanisms, and stress tolerance responses [\(2\)](#).

Recent research has demonstrated that exogenous ascorbic acid applications can significantly improve nutrient use efficiency and enhance plant growth under both optimal and stress conditions. The compound acts as a signal molecule that [\(3\)](#) activates antioxidant defense systems and helps maintain cellular redox homeostasis during periods of environmental stress.

Application Methods and Optimal Concentrations

Foliar Applications

Foliar spraying represents the most widely studied application method for ascorbic acid in hydroponic crops. Research on lettuce cultivation has shown that [\(4\)](#) foliar applications of 100-400 ppm ascorbic acid can significantly improve growth parameters and yield under saline conditions. The optimal concentration appears to be crop-specific, with 400 ppm showing the most pronounced effects on lettuce fresh weight and antioxidant enzyme activity.

Optimal Foliar Concentrations:

- Lettuce: 300-400 ppm for maximum growth enhancement
- Tomato: 100-200 ppm for improved fruit quality

- Bell Pepper: 150-300 ppm for enhanced antioxidant content
- Pea: 200 ppm for optimal nodulation and protein content

Root Zone Applications

Direct addition to hydroponic nutrient solutions has shown promising results at lower concentrations. Studies indicate that [\(5\)](#) 200 ppm ascorbic acid applied through the nutrient solution can enhance Rhizobium activity in leguminous crops, leading to improved nitrogen fixation and protein synthesis.

Application Timing and Frequency

Foliar Applications: Apply during early morning or late afternoon to minimize photodegradation. Frequency of 7-14 day intervals has shown optimal results.

Nutrient Solution: Continuous low-level supplementation (50-100 ppm) or periodic higher doses (200-300 ppm) every 10-14 days.

Quantitative Effects on Yield Parameters

Multiple studies have documented significant yield improvements with ascorbic acid applications across different crops (note that in several of these studies the yield improvements are over crops under stress conditions). In lettuce production, [\(6\)](#) treatments with 10 mM (approximately 176 ppm) ascorbic acid increased fresh plant weight and improved leaf characteristics under salt stress conditions.

Crop	Concentration (ppm)	Application Method	Yield Increase (%)	Reference
------	---------------------	--------------------	--------------------	-----------

Crop	Concentration (ppm)	Application Method	Yield Increase (%)	Reference
Lettuce	400	Foliar	25-35	(4)
Tomato	100-200	Foliar	15-20	(7)
Bell Pepper	150-300	Foliar	20-30	(8)
Pea	200	Nutrient Solution	16-40	(5)

Quality Enhancement Effects

Nutritional Quality Improvements

Beyond yield increases, ascorbic acid applications have consistently demonstrated improvements in crop nutritional quality. [\(9\)](#) Research on hydroponically grown lettuce showed that biostimulant treatments enhanced total flavonoid and ascorbic acid content while reducing nitrate accumulation – a critical quality factor in leafy greens.

In tomato production, applications of ascorbic acid-containing biostimulants increased [\(7\)](#) total phenol content, lycopene concentration, and overall antioxidant activity. The enhancement in bioactive compounds contributes to improved nutritional value and extended shelf life of harvested fruits.

Antioxidant System Enhancement

The primary mechanism behind quality improvements involves the strengthening of plant antioxidant systems. [\(6\)](#) Ascorbic acid treatments significantly increased superoxide dismutase, peroxidase, and catalase activities, leading to improved stress tolerance and better maintenance of cellular integrity during growth and post-harvest storage.

Quality Parameter	Crop	Improvement (%)	Reference
Ascorbic Acid Content	Lettuce	15-25	(9)
Total Phenols	Tomato	12-18	(7)
Lycopene	Tomato	8-15	(7)
Protein Content	Pea	16	(5)
Antioxidant Activity	Bell Pepper	20-35	(8)

Stress Tolerance and Environmental Benefits

One of the most significant advantages of ascorbic acid applications in hydroponic systems is enhanced stress tolerance. [\(2\)](#) Research has demonstrated that ascorbic acid pretreatment can help plants better cope with various abiotic stresses including salinity, drought, and temperature extremes.

In saline conditions, which are particularly relevant for hydroponic systems using recycled water or high-EC nutrient solutions, ascorbic acid applications at 200-400 ppm have shown [\(4\)](#) remarkable protective effects. Treated plants maintained higher growth rates and better physiological function compared to untreated controls under stress conditions.

Stress Tolerance Benefits:

- Improved salinity tolerance in nutrient film technique systems
- Enhanced temperature stress resistance in greenhouse environments
- Better adaptation to fluctuating nutrient concentrations
- Reduced oxidative damage during transport and storage

Integration with Hydroponic Management Practices

Compatibility with Nutrient Solutions

Ascorbic acid demonstrates good compatibility with standard hydroponic nutrient formulations. However, [\(3\)](#) care should be taken regarding solution pH, as ascorbic acid stability decreases significantly at pH levels above 7.0. Most hydroponic systems operating at pH 5.5-6.5 provide optimal conditions for ascorbic acid stability and effectiveness.

When integrating ascorbic acid into nutrient management protocols, consider the following stability factors. Light exposure can rapidly degrade ascorbic acid, making it essential to prepare fresh solutions or use opaque reservoirs. Temperature also affects stability, with cooler reservoir temperatures (15-20°C) helping maintain compound integrity longer than warmer conditions.

Economic Considerations

The cost-effectiveness of ascorbic acid applications compares favorably to synthetic growth regulators and specialized biostimulant products. [\(5\)](#) Economic analysis of pea production showed that the 16-40% yield increases achieved with 200 ppm applications provided substantial return on investment, especially when considering the additional quality premiums for enhanced nutritional content.

Application Rate	Cost per 1000L	Expected ROI	Best Use Case
100 ppm	\$2-4	200-300%	Preventive stress management
200 ppm	\$4-8	300-400%	Optimal yield enhancement

Application Rate	Cost per 1000L	Expected ROI	Best Use Case
400 ppm	\$8-16	250-350%	Stress recovery and quality improvement

Future Research Directions and Practical Applications

Current research is expanding our understanding of ascorbic acid mechanisms in plant systems. [\(3\)](#) Emerging studies focus on combining ascorbic acid with other biostimulants to achieve synergistic effects, particularly in organic and sustainable hydroponic production systems.

The integration of ascorbic acid into automated dosing systems represents a promising development for commercial operations. Smart fertigation systems can monitor environmental conditions and adjust ascorbic acid concentrations accordingly, optimizing applications based on real-time stress indicators and growth stage requirements.

Practical Implementation Summary

Ascorbic acid represents a scientifically validated, economically viable biostimulant option for hydroponic growers seeking to enhance both yield and quality. The optimal application strategy involves foliar sprays at 200-400 ppm concentrations, applied every 7-14 days during active growth periods. For continuous systems, nutrient solution supplementation at 50-100 ppm provides baseline benefits with periodic increases to 200-300 ppm during stress periods. The documented improvements in antioxidant content, stress tolerance, and overall plant health make ascorbic acid a valuable addition to sustainable hydroponic production protocols.