NIR Devices for Leaf Tissue Mineral Analysis

Traditional leaf tissue analysis can cost 50 to 150 USD per sample and take weeks to complete, forcing growers to make nutrient decisions based on outdated information. Near infrared (NIR) spectroscopy devices could theoretically change this reality by providing real time, on site mineral analysis of leaf tissues at a fraction of the cost and time required by conventional laboratory methods.




Image showing NIR measured Vs predicted N values for potatoes, taken from (7)

The Science Behind NIR Technology

Near infrared spectroscopy operates in the electromagnetic spectrum between 700 and 2500 nanometers, measuring the absorption of light by molecular bonds in plant tissues. The technique works by exploiting the fact that organic compounds

containing carbon hydrogen (C-H), oxygen hydrogen (O-H), and nitrogen hydrogen (N-H) bonds absorb specific wavelengths of NIR light (1).

The fundamental principle relies on the relationship between chemical composition and spectral signatures. When NIR light penetrates leaf tissue, different molecules absorb energy at characteristic wavelengths, creating a unique spectral fingerprint. Mathematical models, typically using partial least squares regression (PLSR), then correlate these spectral patterns with actual mineral concentrations determined through traditional analytical methods (2).

Importantly, NIR technology detects macronutrients like nitrogen, phosphorus, and sulfur directly because they are major constituents of NIR sensitive organic compounds such as proteins, nucleic acids, and amino acids. In contrast, nutrients that exist primarily in inorganic forms like calcium, magnesium, and potassium are detected indirectly through their associations with organic compounds (3).

Expected Accuracy Levels

Recent studies show that NIR spectroscopy can achieve excellent prediction accuracy for macronutrients, with correlation coefficients (R^2) typically ranging from 0.80 to 0.95 for nitrogen, phosphorus, and potassium in various crop species (4). Micronutrients generally show lower accuracy, with R^2 values between 0.60 to 0.85, due to their lower concentrations and weaker correlations with NIR active organic compounds.

The ratio of performance to deviation (RPD) values provide another measure of model reliability. RPD values above 2.0 indicate good to excellent predictions, while values above 3.0 are considered excellent for analytical purposes (5). Most successful NIR calibrations for major nutrients achieve RPD values between 2.5 and 4.0, making them suitable for practical

nutrient management decisions.

However, accuracy varies significantly based on sample preparation and measurement conditions. Dried and ground leaf samples consistently produce better calibrations compared to fresh leaves, with improvements in R^2 values of 0.10 to 0.20 for most nutrients. This standardization eliminates moisture content variability and particle size effects that can interfere with spectral measurements (6).

Calibration Challenges and Requirements

Developing robust NIR calibrations requires extensive datasets spanning the full range of nutrient concentrations likely to be encountered in practice. Most successful models require 100 to 300 calibration samples representing different varieties, growth conditions, and nutritional states. The quality of reference analytical data used for calibration directly impacts the final model accuracy, making precise laboratory analysis of training samples essential.

Spectral preprocessing represents another critical calibration challenge. Raw NIR spectra contain noise from light scattering, baseline shifts, and instrument variability that must be corrected before model development. Common preprocessing methods include multiplicative scatter correction (MSC), standard normal variate (SNV), and various derivative transformations, with the optimal approach varying by crop species and nutrient (7).

Model transferability between different instruments, locations, and time periods poses ongoing challenges. Calibrations developed for one NIR device often require recalibration when applied to different instruments, even from the same manufacturer. This limitation necessitates either standardization procedures or the development of universal

calibration models that work across multiple platforms.

Real World Application Issues

Field deployment of NIR devices introduces additional complications not encountered in laboratory settings. Temperature variations can significantly affect spectral measurements, as changing temperatures alter the abundance of organic compounds in plant tissues and the optical properties of the instrument itself (8).

Moisture content represents perhaps the most significant challenge for in field NIR analysis. Water absorption bands can overwhelm nutrient signals in fresh leaf tissue, reducing prediction accuracy by 20 to 40% compared to dried samples. Some portable NIR devices attempt to compensate through moisture correction algorithms, but these approaches add complexity and potential error sources.

Plant species specificity also limits practical implementation. Most NIR calibrations work best for the specific crop and varieties used in model development. Attempting to apply potato leaf calibrations to tomato plants, for example, typically results in poor accuracy. This specificity requirement means that commercial operations need either species specific calibrations or must accept reduced accuracy when using general purpose models.

Comparison with Traditional Analytical Techniques

Parameter	NIR Spectroscopy	ICP-OES	Atomic Absorption	Ion Chromatography
Analysis Time	30 seconds	5-10 minutes per sample	2-5 minutes per element	15-30 minutes

Parameter	NIR Spectroscopy	ICP-OES	Atomic Absorption	Ion Chromatography
Sample Preparation	Minimal (grinding optional)	Acid digestion required	Acid digestion required	Water extraction
Cost per Analysis	\$1-5	\$25-50	\$15-30	\$20-40
Multi-element Capability	Yes (simultaneous)	Yes (simultaneous)	No (single element)	Limited
Accuracy (under ideal calibration and sampling conditions)	Moderate (R ² 0.80-0.95 majors) Poor (R ² < 0.6-0.85 micros)	Excellent (R²>0.99)	Excellent (R²>0.99)	Very Good (R²>0.95)
Detection Limits	Moderate (0.1-1.0%)	Excellent (ppm level)	Very Good (ppm level)	Good (10-100 ppm)
Equipment Cost	\$15,000-50,000	\$150,000-300,000	\$25,000-75,000	\$50,000-100,000
Portability	High (handheld available)	None (lab only)	Low (benchtop)	Low (benchtop)
Chemical Safety	None (no chemicals)	High risk (acids)	High risk (acids)	Low risk
Operator Training	Minimal	Extensive	Moderate	Moderate

Economic Considerations for Commercial Growers

The economics of NIR technology become compelling for operations analyzing more than 200 leaf samples annually. Traditional laboratory analysis costs typically range from 50 to 150 USD per sample including shipping and handling, while NIR analysis costs drop to 1 to 5 USD per sample after initial equipment investment. For a medium scale greenhouse operation testing weekly throughout the growing season, this represents potential savings of 10,000 to 30,000 USD annually.

However, the initial capital investment for quality NIR

equipment ranges from 15,000 to 50,000 USD, depending on spectral range and measurement capabilities. Handheld devices suitable for basic macronutrient analysis start around 15,000 USD, while benchtop instruments capable of full spectrum analysis and micronutrient detection can exceed 50,000 USD (9).

Current Limitations and Future Prospects

Despite significant advances, NIR technology for leaf analysis still faces several limitations. Micronutrient detection remains challenging due to low concentrations and weak spectral signatures. Reliable calibrations for elements like iron, zinc, and manganese typically require concentrations above 100 mg/kg, limiting utility for detecting subtle deficiencies (10).

The development of machine learning approaches and artificial neural networks shows promise for improving prediction accuracy and handling complex spectral relationships. These advanced mathematical techniques can potentially extract more information from NIR spectra than traditional regression methods, particularly for challenging nutrients and mixed species applications. However the success of these techniques hinges on the amount of available data, if the learning library is not big enough, or your crop deviates substantially from it, your accuracy could be even worse than without these complex approaches.

Practical Recommendations

For commercial growers considering NIR technology, the decision should be based on sample volume, required accuracy, and available budget. Operations analyzing fewer than 100 samples annually are generally better served by traditional

laboratory analysis. However, high throughput operations, research facilities, and precision agriculture applications can achieve significant benefits from a well calibrated NIR implementation.

When implementing NIR technology, invest in proper calibration development using samples from your specific crops and growing conditions. Generic calibrations provided by instrument manufacturers rarely achieve the accuracy needed for reliable nutrient management decisions. Plan for ongoing calibration maintenance and periodic validation against traditional analytical methods to ensure continued accuracy. NIR instruments that cannot be properly calibrated for the exact conditions of the grower are much more likely to lead to unusable results.

The future of leaf tissue analysis clearly points toward rapid, non destructive technologies like NIR spectroscopy. While current limitations prevent complete replacement of traditional methods, NIR devices offer valuable screening capabilities and real time insights that can significantly improve nutrient management efficiency under ideal conditions. As the technology continues to mature and costs decrease, adoption will likely accelerate across all scales of agricultural production.

Oxygenation of Nutrient Reservoirs in Substrate-Based Soilless Crops

When growers new to hydroponics start working with soilless systems, one of the first questions that comes up is whether

they should be actively aerating their nutrient solutions. Air stones bubbling away in reservoirs have become synonymous with hydroponics, particularly in deep water culture systems. However, when growing in substrates like coconut coir or rockwool, the situation is fundamentally different. Understanding where root oxygen comes from in substrate systems can help you avoid wasting resources on unnecessary equipment while also helping you understand the real limitations of these growing methods.




Figure 1. Influence of particle size on air and water distribution in growing substrates. Coarse particles create macropores that hold air after drainage, while fine particles create micropores that retain water. The balance between these determines oxygen availability to roots $(\frac{7}{2})$.

Where Roots Get Oxygen in Substrate Systems

In substrate-based growing systems, roots obtain nearly all their oxygen from air-filled pores within the growing medium, not from dissolved oxygen in the nutrient solution. Substrates like rockwool and coconut coir typically have total porosities exceeding 80%, compared to typical soil porosities below 40% (1). This high porosity ensures there are enough water-filled pores for nutrient transport as well as enough air-filled pores for oxygen transport.

The key parameter governing oxygen availability in substrates

is air-filled porosity, which represents the percentage of air contained in a fixed volume of substrate after it has been saturated with water and the free water has drained (2). Research on growing media has shown that adequate air-filled porosity levels for optimal plant growth typically range from 10-20%, with some studies suggesting that values above 20% may be necessary immediately after irrigation to prevent hypoxia (3).

When you irrigate a substrate, the nutrient solution displaces air in the open pores. As the substrate drains, air is drawn back down into the root system. This cycle of wetting and drying is what supplies roots with fresh oxygen. The oxygen diffusion coefficient in air is approximately 10,000 times higher than in water, which means that gas-phase oxygen transport through substrate pores is far more efficient than dissolved oxygen transport through water (4).

Substrate Type	Total Porosity (%)	Air-Filled Porosity at Field Capacity (%)	Water Holding Capacity (%)
Rockwool	95-97	15-20	75-80
Coconut Coir	85-90	20-30	60-70
Coco/Perlite (70:30)	85-90	25-35	55-65
Perlite	50-70	30-40	30-40

Does Nutrient Solution Oxygenation Make Sense?

The short answer is that in properly managed substrate systems with adequate irrigation frequency, oxygenating the nutrient solution in your reservoir provides minimal benefit to plant growth. The reason is simple: the overwhelming majority of oxygen uptake occurs through gas-phase diffusion in the airfilled pores of the substrate, not through dissolved oxygen in the water phase.

Research comparing water-based and substrate-based cultivation systems has demonstrated that substrate-grown plants can thrive even when oxygen supply through irrigation is potentially growth limiting, as long as the substrate maintains adequate air-filled porosity (1). In contrast, water culture systems where roots are continuously submerged rely entirely on dissolved oxygen, making aeration critical in those applications.

The irrigation strategy you use has far more impact on root zone oxygen than dissolved oxygen levels in your reservoir. Allowing substrates to dry down between irrigations increases air-filled porosity and draws fresh air into the root zone. Over-irrigation is far more likely to cause oxygen deficiency problems than low dissolved oxygen in your nutrient tank. When substrates remain saturated, air-filled pores fill with water, creating anaerobic conditions regardless of how much you aerate your reservoir.

The exception to this general rule would be in situations where you have continuous or very frequent irrigation with minimal drainage, essentially converting your substrate system into something closer to a water culture system. In such cases, dissolved oxygen becomes more important, but this represents poor management of a substrate system rather than a reason to add aeration.

The Pathogen Risk of Solution Aeration

While aerating nutrient solutions might seem harmless even if unnecessary, there is a significant downside that growers should consider: the increased risk of introducing and spreading waterborne pathogens, particularly species of Pythium and Phytophthora.

These oomycete pathogens are among the most problematic diseases in hydroponic systems. They produce motile zoospores that can swim through nutrient solutions using flagella,

allowing them to spread rapidly through recirculating systems $(\underline{5})$. When closed hydroponic systems are used, pathogens can enter and then rapidly disseminate, particularly during periods of stress such as high temperatures or low dissolved oxygen levels $(\underline{5})$.

Aeration systems create several opportunities for pathogen introduction and proliferation. Air stones and diffusers provide surfaces for biofilm formation where pathogens can colonize. The turbulence created by aeration helps distribute any pathogens present throughout the solution more effectively than they would spread by passive diffusion. The air being pumped into the system can carry airborne pathogen propagules, and unless you are using sterile filtration on your air intake, you are essentially inoculating your reservoir with whatever microorganisms happen to be in your growing environment.

Low dissolved oxygen has been reported to increase Pythium infection in hydroponic systems (6). However, in substrate systems where roots obtain oxygen primarily from air-filled porosity rather than dissolved oxygen, the relationship between solution aeration and disease suppression becomes less clear. The more relevant factors for disease prevention in substrate systems include maintaining proper irrigation frequency to ensure adequate substrate aeration, avoiding prolonged saturation, and keeping solution temperatures below 24°C where practical.

Pathogen Risk Factor	Risk Level with Aeration	Risk Level without Aeration
Airborne contamination introduction	High	Low
Pathogen distribution through solution	High (turbulent mixing)	Moderate (passive diffusion)
Biofilm formation sites	High (air stones, tubing)	Low (tank surfaces only)

Pathogen Risk Factor	Risk Level with Aeration	Risk Level without Aeration
Solution temperature increase	Possible (pump heat)	Minimal

Practical Recommendations


For growers using substrate-based systems, the evidence suggests that resources are better spent on proper irrigation management than on solution aeration. Focus on selecting substrates with adequate air-filled porosity, implementing irrigation schedules that allow periodic drying to refresh the air in the root zone, and maintaining appropriate solution temperatures.

If you are growing in pure water culture systems like deep water culture, NFT, or aeroponics, then maintaining adequate dissolved oxygen becomes critical and aeration or other oxygenation methods are necessary. But if you are growing in rockwool, coco coir, or similar substrates with good drainage, your plants are getting their oxygen from the air in the substrate pores, not from the water in your reservoir.

The key takeaway is this: in substrate systems, oxygen management happens at the substrate level through proper irrigation practices, not at the reservoir level through aeration. Understanding this fundamental difference can help you avoid unnecessary equipment costs while potentially reducing your risk of introducing waterborne pathogens into your growing system.

Top 5 Open Source Hardware Tools to Boost Your Hydroponic Yields

The equipment you use to monitor and control your hydroponic system can make or break your crop quality and yields. Commercial systems often cost thousands of dollars, putting precision agriculture out of reach for small to medium-scale operations. Fortunately, open source hardware platforms like Raspberry Pi and Arduino have revolutionized what growers can achieve with limited budgets. In this post, I'll walk you through five open source hardware tools you can build yourself to significantly improve your operation.



Results of the plant phenotyping system used in (4)

Why Open Source Hardware Matters

Before diving into specific tools, it's worth understanding why open source hardware is such a game-changer for hydroponics. These platforms typically cost 90-95% less than commercial equivalents while offering comparable or superior functionality (1). More importantly, you control the design, can modify it for your specific needs, and aren't locked into proprietary systems.

Automated pH and EC Control System

Maintaining optimal pH (typically 5.8-6.2) and electrical conductivity is critical for nutrient uptake and plant health. Manual adjustment is time-consuming and prone to error. An automated system using a Raspberry Pi 3 with fuzzy logic control can maintain these parameters with remarkable precision (2).

The system uses pH and EC sensors as inputs and controls four pumps (high EC solution, water, acid, and base) to automatically adjust your nutrient solution. In controlled trials with lettuce, automated systems maintained target pH within 0.04 units and achieved 7% greater leaf width compared to manual management (2). The entire system can be built for under \$200 using readily available components.

Key Benefits:

Parameter	Improvement
pH stability	±0.04 units
Labor reduction	90% reduction in manual testing
Plant growth	7% increase in leaf width
System cost	~\$150-200

2. Open Source PAR Sensor

Light is your most important input for photosynthesis, yet many growers operate blindly without measuring photosynthetically active radiation (PAR). Commercial PAR sensors cost \$600-1000, but you can build an open source version using an AS7341 spectral sensor and ESP32 microcontroller for approximately \$50-70 (1).

This system measures PAR across the 400-700nm spectrum with accuracy comparable to commercial sensors (within 2-5% error). It includes data logging to an SD card and optional WiFi connectivity for remote monitoring. Since a 1% increase in lighting typically provides a 1% increase in yield (3), knowing exactly how much light your plants receive allows you to optimize your lighting strategy and maximize productivity.

3. Plant Phenotyping Camera System

Visual monitoring of plant growth provides invaluable data for optimizing your system. A Raspberry Pi-powered imaging system can capture time-lapse images of your crops and extract quantifiable traits like plant area, height, and color (4).

For around \$100 per camera unit, you can set up multiple Raspberry Pi cameras in your grow space to continuously monitor plant development. The images can be processed using open source software like PlantCV to automatically measure growth rates, detect stress before it's visible to the naked eye, and compare different treatments or varieties (4). This approach scales well, with some research groups successfully deploying 180 cameras to monitor 1800 plants simultaneously with 96% uptime (5).

Phenotyping System Comparison:

System Type	Cost per Unit	Traits Measured	Temporal Resolution
Commercial	\$5,000-50,000	Multiple	Variable
Raspberry Pi	\$100-150	Area, height, color, shape	Minutes to hours
Manual	Labor cost	Limited	Daily at best

4. Multi-Parameter Environmental Data Logger

Environmental conditions directly impact both yield and quality. An Arduino or ESP32-based data logger can simultaneously monitor temperature, humidity, CO2, dissolved oxygen, and solution temperature. By logging data every 1-2 minutes, you can identify patterns and problems that periodic manual measurements would miss (6).

Dissolved oxygen is particularly important but rarely monitored by small-scale growers due to sensor costs. However, maintaining adequate dissolved oxygen levels (above 5 mg/L) is essential for root health and nutrient uptake. An open source system using Atlas Scientific sensors can monitor DO along with other parameters for a few hundred dollars.

The real value comes from the data. When you can correlate environmental conditions with plant performance, you can make informed decisions about climate control, identify the optimal conditions for your specific varieties, and catch problems before they impact yields.

5. Automated Nutrient Dosing System

Precision in nutrient delivery improves both crop quality and reduces waste. An automated dosing system using peristaltic pumps controlled by a microcontroller can deliver exact amounts of different nutrient solutions based on real-time sensor feedback (2).

The system can be programmed to maintain target EC and pH values by calculating the required pump run times using mathematical models or machine learning approaches. More sophisticated implementations can adjust nutrient ratios based on plant growth stage or environmental conditions. While commercial fertigation systems cost thousands of dollars, an open source version can be built for \$300-500 depending on the number of nutrient solutions you want to control.

Getting Started

Building these systems requires some technical knowledge, but the barrier to entry is lower than you might think. Start with a single-purpose system like the PAR sensor or environmental data logger to learn the basics. Online communities around Arduino, Raspberry Pi, and platforms like Mycodo provide extensive documentation and support.

The investment in time to build and configure these systems pays dividends through improved crop quality, higher yields, and better understanding of your growing environment. Even if you only implement one or two of these tools, you'll gain capabilities that were reserved for large commercial operations just a few years ago.

Cost Comparison:

Tool	Commercial Cost	Open Source Cost	Savings
pH/EC Controller	\$800-2000	\$150-200	85-90%
PAR Sensor	\$600-1000	\$50-70	93-95%
Phenotyping System	\$5000+	\$100-150	97%
Data Logger	\$400-800	\$100-200	60-80%
Dosing System	\$1500-3000	\$300-500	75-85%

The beauty of open source hardware is that you can start small and expand as your needs grow. Each tool you add gives you more control and insight into your operation, ultimately leading to better crops and more profitable growing.

Have you built any open source monitoring or control systems for your hydroponic operation? What challenges did you face and what benefits have you seen? Let me know in the comments below!

Growing Soilless Crops Without Nitrates: Practical Options When Nitrate Salts Are Unavailable

For growers in regions where geopolitical conflicts or economic constraints limit access to nitrate fertilizers like calcium nitrate and potassium nitrate, the question arises: can you grow hydroponic or soilless crops using only alternative nitrogen sources? The short answer is yes, but with important limitations and necessary substrate modifications. This post explores the science behind nitrate-free soilless growing and practical strategies for growers facing nitrate scarcity.



Figure 1. Effects of nitrate concentration (25, 50, 75, 100 and 150% of the recommended dose) and proportion of nitrate/ammonium (0:100, 25:75, 50:50, 75:25 and 100/0) in the nutrient solution for hydroponics, on the development of lettuce Iceberg type.

The above image is sourced from (8).

Why Nitrates Dominate in Hydroponics

In conventional hydroponics, 85-95% of nitrogen is supplied as nitrate (NO3-) rather than ammonium (NH4+). This preference exists for good reasons. Plants can safely store nitrate in vacuoles without toxicity, while ammonium accumulation in plant tissues causes rapid damage (1). In soil, nitrifying bacteria convert ammonium to nitrate before plant uptake, but most soilless substrates lack these microbial communities. Without this conversion, ammonium concentrations that would be harmless in soil become highly toxic in hydroponics.

Research on tomatoes shows that plants supplied with 112 ppm nitrogen as ammonium developed severe toxicity symptoms and produced only one-third the biomass of nitrate-fed plants (1). Even at 14 ppm nitrogen, ammonium-only nutrition suppressed growth compared to mixed nitrogen sources. For lettuce, similar effects occur, with crown discoloration and biomass

reductions appearing at 50 ppm ammonium nitrogen (2).

Maximum Safe Ammonium Levels

The tolerance threshold varies by species and conditions, but general guidelines exist:

Crop Type	Maximum Safe Ammonium (% of total N)	Maximum Concentration (ppm N)
Most crops (standard)	10-15%	15-30 ppm
Sensitive crops (tomato, pepper, lettuce)	5-10%	10-20 ppm
Cold conditions (<15°C)	0 - 5%	0-10 ppm
High light, fast growth	15-20%	20-40 ppm

These limits exist because ammonium uptake is passive and rapid, plants cannot regulate it effectively, and it disrupts calcium and magnesium uptake while acidifying the root zone (3).

Substrate Amendments: Creating Artificial Soil

The key to using higher ammonium levels or organic nitrogen sources is establishing nitrifying bacteria in the substrate. Recent research demonstrates that soilless substrates can be inoculated with microbial communities that convert organic nitrogen to nitrate (4).

Effective substrates for nitrification include rockwool, vermiculite, polyurethane foam, oyster shell lime, and rice husk charcoal. The process requires:

1. Inoculum source: Bark compost or mature vermicompost

provides ammonia-oxidizing bacteria (AOB) and nitrite-oxidizing bacteria (NOB). Add 1g per 100mL substrate initially.

- 2. **Temperature**: Nitrifying bacteria function optimally at 25-42°C. Below 15°C, nitrification slows dramatically, causing ammonium accumulation (5).
- 3. **Humidity and aeration**: Substrates need >50% relative humidity and adequate oxygen. Waterlogged conditions inhibit nitrification and promote denitrification.
- 4. **Establishment period**: Allow 2-3 weeks for bacterial colonization before planting. Daily additions of dilute organic fertilizer (6 mg N per 100mL substrate) accelerate establishment.

Practical Nitrogen Sources

Ammonium Salts

Ammonium sulfate ((NH4)2S04) is the most accessible ammonium source globally. At 21% nitrogen, it provides both N and sulfur. However, use caution:

- Never exceed 20% of total nitrogen as ammonium in solution
- Monitor substrate pH closely, as ammonium uptake releases protons and acidifies the root zone
- Increase ratios only under high light and warm temperatures (>20°C)
- Sensitive crops like lettuce, tomato, and pepper tolerate lower ratios

Ammonium phosphate (MAP or DAP) offers nitrogen plus phosphorus but requires even more careful management due to rapid pH shifts.

Urea

Urea (CO(NH2)2) at 46% nitrogen is economical and widely available. In water, urease enzymes (either from bacteria or added exogenously) hydrolyze urea to ammonium. However, hydroponic studies on various crops show that urea performs poorly as a sole nitrogen source (6). Plants fed only urea exhibited nitrogen deficiency symptoms at low concentrations and toxicity at high concentrations. The primary issues are:

- Insufficient uptake of intact urea by most crop species
- Variable conversion rates without soil bacteria
- pH instability during hydrolysis

Combined applications of urea with nitrate showed better results than urea alone, but if nitrates are unavailable, urea offers limited benefit beyond what ammonium salts provide (6).

Compost and Organic Extracts

Compost leachates and vermicompost teas contain nitrogen primarily as proteins, amino acids, and ammonium. Direct use in inert hydroponics fails because plants cannot efficiently absorb complex organic nitrogen. However, two approaches work:

Aerobic nitrification method: Add organic nitrogen sources like corn steep liquor (1g/L) or fish emulsion plus bark compost (0.5g/L) as bacterial inoculum. Aerate for 12 days, during which bacteria convert organic N and ammonium to nitrate, reaching 100-130 ppm N as nitrate (7). This creates a low-cost, nitrate-containing solution from readily available materials.

Substrate-based mineralization: Inoculate substrates with compost microbes and apply dilute organic fertilizers daily. The substrate acts as a biofilter, mineralizing organic N to nitrate before plant uptake (4). This method requires 2-3

weeks establishment and careful moisture management.

Expected Yield Impacts

When managed properly with substrate amendments and bacterial communities, yields can approach conventional hydroponic levels. Studies show that tomatoes grown with nitrified organic solutions performed comparably to mineral fertilizer controls when adequate nitrate was generated (7).

However, several factors reduce yields in poorly managed nitrate-free systems:

- Ammonium toxicity: High ammonium causes 30-70% yield reductions across most crops (1)
- **Nutrient imbalances**: Ammonium competes with Ca²⁺ and Mg²⁺ uptake, inducing deficiencies
- pH instability: Root zone acidification from ammonium uptake reduces nutrient availability
- Incomplete mineralization: Organic N sources may not fully convert to plant-available forms

Realistic expectations for growers transitioning to nitratefree systems:

- First crop cycle: 50-70% of conventional yields while optimizing conditions
- Established systems with functioning bacterial communities: 80-95% of conventional yields
- Cold season growing (<15°C): 40-60% due to impaired nitrification

Nutrient Solution Modifications

Without calcium nitrate, calcium must come from chloride or

sulfate sources rather than nitrate. Calcium chloride is highly soluble but adds chloride. Gypsum (calcium sulfate) doesn't have the solubility needed to make concentrated stock solutions and therefore can only be added to the final solutions or added to the media as an amendment. Calcium chloride can add unwanted high amounts of chlorides as it's therefore best avoided. If you are doing composting amendments then limestone amendments might be the most desirable way to supply Ca to the crop.

Critical Success Factors

To successfully grow soilless crops without nitrate fertilizers:

- 1. **Establish nitrifying bacteria**: This is non-negotiable for using organic N or high ammonium levels
- 2. **Monitor pH constantly**: Ammonium acidifies solutions; maintain pH 5.8-6.5 through buffering or base addition
- 3. **Provide adequate calcium**: Use calcium chloride or sulfate since calcium nitrate is unavailable
- 4. Keep temperatures warm: >20°C substrate temperature for bacterial activity
- 5. **Start conservatively**: Begin with 10% ammonium and increase gradually as plants adapt
- 6. Choose tolerant species first: Leafy greens like pak choi are more tolerant than tomatoes or peppers

Conclusion

Growing soilless crops without nitrates is achievable but requires different management than conventional hydroponics. The approach depends on creating conditions that mimic soil processes, establishing microbial communities to convert ammonium and organic nitrogen to nitrate within the substrate. While yields may initially be lower, proper substrate

inoculation, temperature management, and careful nitrogen source selection can produce acceptable results. For growers with limited access to nitrate salts, combining small amounts of ammonium sulfate (20-30 ppm N) with aerobically nitrified compost teas or inoculated substrates offers the most practical path forward.

Using Portable Low-Cost Chlorophyll Sensors to Assess Plant Health and Improve Crop Quality in Hydroponics

When you grow plants hydroponically you become responsible for delivering the exact amount of every essential nutrient. Getting nitrogen right is particularly challenging since plants can require dramatically different amounts depending on their growth stage. Traditional methods to assess nitrogen status require sending leaf samples to a lab, which is expensive, destructive, and provides results too late to make timely corrections. Portable chlorophyll meters offer a practical solution.



A DIY chlorophyl meter compared to some commercial alternatives (taken from (9)).

What Are Portable Chlorophyll Meters?

Portable chlorophyll meters are handheld devices that non-destructively estimate the chlorophyll content in plant leaves. The most widely used device is the SPAD-502 meter, which works by measuring light transmission through a leaf at two wavelengths: 650 nm (red light, which chlorophyll absorbs) and 940 nm (infrared, which chlorophyll does not absorb). The device calculates a dimensionless SPAD value based on the transmission ratio (1). Since 50-70% of leaf nitrogen is contained in chlorophyll molecules, these readings provide a reliable proxy for nitrogen status (2).

These meters are particularly useful in hydroponic systems where you have complete control over nutrient delivery and can make rapid adjustments when deficiencies are detected. Research has demonstrated strong correlations between

chlorophyll meter readings and nitrogen status in major hydroponic crops including tomato (3), lettuce (4), and greenhouse vegetables (5).

Major Advantages

The primary advantage is that measurements are instantaneous and non-destructive. You can measure the same leaf repeatedly throughout the growing season without harming the plant. This is especially valuable in hydroponics where you might want to monitor nitrogen status weekly or even daily during critical growth periods.

The correlation between SPAD readings and leaf nitrogen concentration is typically very strong. In romaine lettuce grown in soilless culture, SPAD readings showed correlation coefficients of $R^2 = 0.90$ with nitrogen concentration and $R^2 = 0.97$ with chlorophyll content (4). Similar results have been reported for greenhouse tomatoes, where R^2 values ranged from 0.86 to 0.94 (6).

Unlike laboratory analysis, chlorophyll meters provide immediate feedback. When you detect that SPAD readings are dropping below your target range, you can adjust your nutrient solution that same day, particularly advantageous in fertigation systems (7).

Low-Cost Alternatives

The SPAD-502 meter typically costs \$2,000-\$2,600, which can be prohibitive for small growers. Several low-cost alternatives have been developed and validated. The atLEAF meter costs around \$200-\$300 while providing equivalent performance (8). Studies found strong correlations ($R^2 = 0.96$) between SPAD and atLEAF meters across multiple crop species (8).

Functional chlorophyll meters can even be built from scratch

using simple electronic components for under \$100. A recent study described construction using 3D-printed hardware and off-the-shelf LEDs and photodiodes that achieved strong correlations with both the SPAD-502 and atLEAF meters (9).

Device	Cost (USD)	Wavelengths (nm)	Key Features
SPAD-502	2,000-2,600	650, 940	Industry standard
atLEAF+	200-300	660, 940	Data logging, SPAD conversion
MC - 100	400-600	653, 931	Larger measurement area
Custom Arduino	<100	650, 940	Requires assembly

The atLEAF meter is available through <u>agricultural supply</u> <u>retailers</u>, while various manufacturers offer devices in the \$100-\$300 range through online platforms.

Research in Hydroponic Crops

Chlorophyll meters have been successfully used to guide nitrogen management in various hydroponic crops. In tomato production, using SPAD readings to trigger nitrogen applications resulted in the highest yields compared to fixed-rate applications, with improved nitrogen use efficiency (3). Researchers established critical SPAD values for different physiological stages, allowing growers to apply nitrogen only when needed.

For lettuce grown in high tunnels with fertigation, both SPAD and atLEAF meters accurately estimated nitrogen status, fresh weight, and chlorophyll concentration with R^2 values above 0.90 (4). Research on basil with different nitrogen rates showed that SPAD, atLEAF, and MC-100 meters all provided reliable estimates with R^2 values of 0.93-0.98 (10).

Important Limitations

While valuable, chlorophyll meters have limitations growers need to understand. The relationship between SPAD readings and actual nitrogen content can vary between species. Research on seven crop species found that while the relationship between SPAD and chlorophyll content was consistent, the relationship between SPAD and leaf nitrogen varied widely (1). You cannot use the same threshold values across different crops.

Environmental factors also affect readings:

- **Time of day**: Chloroplast movement can cause SPAD readings to decrease by 13-28% at midday under low nitrogen conditions (1). Take readings in early morning or maintain consistent measurement times.
- **Light history**: Short-term changes in growth light affect nitrogen allocation to chlorophyll (1).
- Leaf position and age: Chlorophyll content varies across leaf positions and with age. Always measure the same leaf position.

Chlorophyll meters provide relative rather than absolute measurements. To use them effectively, you need to establish calibration curves for your specific crop and growing conditions.

Best Practices

Establish Baseline Values: Grow plants at different nitrogen levels and measure both SPAD readings and leaf nitrogen concentration via lab analysis. This establishes your calibration curve.

Use Reference Strips: Maintain a section of plants receiving optimal nitrogen. Compare readings from your bulk crop to these reference plants. If bulk readings drop more than 5-10%

below reference, increase nitrogen delivery (7).

Standardize Protocol: Always measure the same leaf position. For leafy greens, measure the most recently fully expanded leaf. For tomatoes, measure leaflets on the leaf closest to the most recent fruit cluster. Take measurements at the same time daily, preferably early morning (5).

Take Multiple Readings: SPAD readings can vary 10-15% between individual plants. Measure at least 20-30 plants per zone and use the average (7).

Species-Specific Calibration: If you grow multiple crops, establish separate calibration curves for each.

Crop	Optimal Range	Action Threshold	Measurement Location
Lettuce	35-45	<32	Youngest fully expanded leaf
Tomato	45-55	<42	Leaflet near newest fruit cluster
Cucumber	40-50	<37	3rd fully expanded leaf
Basil	35-45	<32	Terminal leaves

Note: These are general guidelines. Establish specific thresholds for your cultivars through calibration with lab analysis.

Conclusions

Portable chlorophyll meters represent an excellent investment for hydroponic growers optimizing nitrogen management. Low-cost alternatives make this technology accessible even for hobby growers. While these devices have limitations related to species-specific calibration and environmental factors, following standardized protocols allows effective use for management decisions.

The key is understanding that chlorophyll meters provide relative measurements. Take time to establish proper baseline values for your crops and conditions. Once calibrated, these devices help fine-tune nitrogen delivery, reduce fertilizer waste, prevent deficiencies, and improve crop yield and quality.

For growers ready to adopt this technology, starting with an <a href="https://attended.com/atten

The Problems with Brix Analysis of Sap in Crops

Brix analysis, the measurement of soluble solids in plant sap using a refractometer, has gained popularity as a quick field test for assessing plant health and crop quality. The method is appealingly simple: squeeze some sap from a leaf onto a refractometer, and within seconds you get a number that supposedly tells you how healthy your plant is. Many proponents claim that plants with high brix readings are more resistant to pests and diseases, while low readings indicate nutritional problems. However, when we examine the scientific literature surrounding brix measurements in plant sap, particularly for agronomically important crops in hydroponic or soilless systems, we find that this technique has substantial limitations that are often overlooked.



A refractometer, the most common tool to measure brix of plant sap.

The Appeal and the Theory

The basic premise of brix analysis is straightforward. The refractometer measures the refractive index of a solution, which correlates with the concentration of dissolved solids (1). In plant sap, these dissolved solids include sugars, amino acids, proteins, minerals, and other organic compounds. The theory suggests that healthier plants with better nutrition will have higher sugar content from improved photosynthesis, leading to higher brix readings (2). While this sounds reasonable, the reality is far more complex.

Problem 1: Dramatic Diurnal Variation

One of the most significant issues with brix measurements is their extreme variability throughout the day. Plants accumulate sugars during photosynthesis in the light period and then mobilize these sugars at night for growth, respiration, and transport to sink organs. Research on mature oak trees showed that total leaf sugars increased by an average of 16 mg/g dry weight during the day and returned to baseline at night (2). This represents substantial diurnal fluctuation that can produce 30% or more variation in brix readings depending on time of day (3).

Studies on maize have shown that starch and soluble sugars in leaves follow predictable diurnal patterns, with soluble carbohydrates peaking in the afternoon and reaching their minimum before dawn (3). The timing of peak brix values varies by species and growing conditions. Some plants show maximum sugar accumulation at midday, while others peak in the afternoon (2). This means that a brix reading taken at 10 AM might be dramatically different from one taken at 3 PM on the same plant, even though the plant's nutritional status has not changed.

Weather conditions further complicate matters. Plants have been observed to move sugars to roots in anticipation of storms, sometimes days in advance, causing brix readings to drop substantially even though the plant is not experiencing nutritional stress. Water stress also affects readings, as dehydration concentrates dissolved solids and artificially elevates brix values without indicating better plant health.

Problem 2: Spatial Variation Within Plants

The location where you sample sap makes an enormous difference in the reading you obtain. Research has consistently shown large differences in sugar content between young and old leaves, with old leaves often having substantially different concentrations than new growth (2). In reproductive plants, leaves near fruits typically show the lowest brix readings because fruits have high nutritional demands and act as strong

sinks for sugars and other nutrients.

This spatial heterogeneity means that two technicians sampling the same plant but choosing different leaves could easily obtain readings that vary by 50-70%. Without strict standardization of which leaf to sample, when during its development, and from which position on the plant, brix measurements become nearly impossible to compare across samples or over time.

The Logistical Challenge

For brix analysis to be useful as a management tool, it requires an extraordinary level of commitment and consistency. You would need to collect samples at different locations on each plant, within different areas of your growing system, under different weather conditions, and critically, at exactly the same time of day, multiple times per week (4). Because of this inherent variability, effective use requires managing trends rather than individual measurements. Most growers simply do not have the bandwidth to develop the degree of familiarity needed with brix readings for it to become a truly reliable diagnostic tool.

What Brix Cannot Tell You

Perhaps most importantly, even if you could control for all the temporal and spatial variation, brix readings provide very limited actionable information. A low brix reading tells you that soluble solids are low at that moment, but it does not tell you why. Is it a nitrogen deficiency? Phosphorus? Calcium? Is it a problem with root function? Temperature? Light intensity? Water relations? The brix value alone provides no way to differentiate between these possibilities.

Additionally, brix measurements tell you nothing about immobile nutrients like calcium and boron, which do not move

readily through the sap. These nutrients are critical for cell wall formation, disease resistance, and fruit quality, yet they remain essentially invisible to brix analysis.

A Better Alternative: Leaf Tissue Analysis

When growers need reliable information about plant nutritional status in hydroponic systems, leaf tissue analysis provides a far more comprehensive and actionable alternative. Unlike brix analysis, which measures only mobile compounds in sap at a single moment, tissue analysis quantifies the total accumulated concentrations of both mobile and immobile nutrients in plant tissues (4).

Tissue analysis provides specific concentration values for nitrogen, phosphorus, potassium, calcium, magnesium, sulfur, and all essential micronutrients. These values can be compared against established sufficiency ranges for your specific crop, allowing you to identify which nutrients are deficient, adequate, or excessive. This specificity enables targeted corrective actions rather than guesswork.

While tissue analysis does require sending samples to a laboratory and waiting for results, it provides a stable measurement that is far less affected by time of day or recent environmental fluctuations. Modern labs can return results within days, and the interpretive frameworks for tissue analysis are well-established across hundreds of crop species.

Comparison Factor	Brix Analysis	Tissue Analysis
Time of day sensitivity	Very high (30-70% variation)	Low
Spatial variation within plant	Very high	Moderate

Comparison Factor	Brix Analysis	Tissue Analysis
Nutrients detected	Soluble solids only (mostly sugars)	All essential elements (15+)
Specificity	Non-specific	Element-specific
Interpretation	Difficult without extensive experience	Well-established sufficiency ranges
Cost per sample	Low	Moderate
Actionable information	Limited	Comprehensive

Practical Recommendations

This is not to say that refractometers have no place in crop monitoring. For specific applications like determining harvest timing for fruits or monitoring sugar accumulation in reproductive organs, brix can be useful. However, for assessing the overall nutritional health of vegetative crops in hydroponic systems, the limitations of sap brix analysis are substantial.

If you are serious about optimizing nutrition in your hydroponic operation, invest in regular tissue analysis rather than relying on brix readings. Sample the most recently matured leaves at consistent growth stages, submit samples to a reputable agricultural laboratory, and use the results to make informed adjustments to your nutrient formulation and delivery. This approach will provide you with reliable, actionable data that can actually improve your crops, rather than numbers that fluctuate wildly based on time of day and sampling location.

The appeal of a quick field test is understandable, but in the case of brix analysis for plant health assessment, the simplicity comes at the cost of reliability and utility.

Sometimes the best tools are not the fastest ones, and when it comes to understanding what your plants need, there is no substitute for comprehensive analysis.

Comparing Nutrient Solutions for Hydroponic Strawberry Production

Getting the right nutrient solution for strawberries in hydroponics can feel like trying to solve a puzzle where every piece matters. Unlike many crops where you can get away with a generic formula, strawberries are particularly responsive to nutrient composition, especially when it comes to the balance between nitrogen and potassium. Today, we will explore how different nutrient formulations affect both yield and fruit quality in soilless strawberry production.



The Modified Steiner Approach

When researchers at the Technological Institute of Torreón tested different nitrogen and potassium combinations in strawberries, they discovered something important about how these two nutrients interact. Using a (1) modified version of Steiner's Universal Nutrient Solution, they evaluated twelve different formulations with nitrogen ranging from 126 to 210 ppm and potassium from 195 to 430 ppm.

The results were revealing. Plants receiving 168 ppm nitrogen combined with 430 ppm potassium achieved yields of 114 grams per plant, which was significantly higher than lower nitrogen treatments. However, here is where it gets interesting: while high nitrogen boosted yield, it actually decreased fruit quality. The highest soluble solids content (10.5 degrees Brix) occurred at the lowest nitrogen level of 126 ppm. This creates a real dilemma for growers who want both high yields and premium quality fruit.

Solution Type	N (ppm)	P (ppm)	K (ppm)	Ca (ppm)	Mg (ppm)	Yield	Quality Impact
Modified Steiner (Low N)	126	46	195	449	121	89.3 g/plant	Highest Brix (10.5°)
Modified Steiner (Medium N)	168	32	273	360	97	108 g/plant	Moderate Brix (10.0°)
Modified Steiner (High N)	210	19	194	413	111	111 g/plant	Lowest Brix (9.5°)

The Critical Role of Potassium

What emerged from this study was potassium's profound impact on fruit quality. When potassium was increased to 430 ppm, the soluble solids climbed to 10.6 degrees Brix, and phenolic compounds reached their peak as well. The (1) research showed that the optimal combination for maximizing both yield and nutraceutical quality was 168 ppm nitrogen with 430 ppm potassium, resulting in antioxidant capacity of 6305 microequivalents of Trolox per 100 grams.

This makes physiological sense. Potassium plays a fundamental role in sugar transport through the phloem, and when potassium availability is adequate, more sugars accumulate in the fruit. Meanwhile, excessive nitrogen tends to promote vegetative growth and the synthesis of nitrogen containing compounds like proteins and amino acids, rather than the accumulation of secondary metabolites that contribute to fruit quality.

Optimizing NPK Ratios for Chinese Greenhouses

A comprehensive study from China Agricultural University took a different approach by examining the combined effects of nitrogen, phosphorus, potassium, and water on strawberry production. Using a (2) quadratic regression design with 36 treatments, researchers determined that nitrogen was by far the most important factor, followed by water, then phosphorus, with potassium having the least impact on the sweetness to acidity ratio.

Their optimal formulation for achieving yields above 110 grams per plant with excellent fruit quality included nitrogen at 156 to 172 ppm (supplied as calcium nitrate), phosphorus at 54 to 63 ppm (as sodium dihydrogen phosphate), and potassium at 484 to 543 ppm (from potassium sulfate). This represents

significantly higher potassium levels than the Steiner based formulations, suggesting that when other nutrients are optimally balanced, strawberries can benefit from even more potassium.

Nutrient	Optimal Range (ppm)	Impact on Yield	Impact on Quality (SSC/TA)
Nitrogen (N)	156 to 172	Most significant positive effect	Most significant factor
Phosphorus (P)	54 to 63	Moderate positive effect	Second most important
Potassium (K)	484 to 543	Significant positive effect	Minimal impact
Water	12.0 to 13.1 L/plant	Second most important	Third most important

The Calcium and Electrical Conductivity Question

While much attention focuses on NPK ratios, calcium concentration matters enormously in strawberry production. In the modified Steiner solutions, calcium ranged from (1) 244 to 449 ppm depending on the treatment. Higher calcium levels corresponded with lower nitrogen and potassium concentrations, maintaining appropriate osmotic potential.

Research has shown that the electrical conductivity (EC) of the nutrient solution significantly impacts strawberry performance in soilless culture. Studies using different EC levels found that (3) 1.3 mS/cm was optimal for spring production, while 2.2 mS/cm proved better during winter months. This seasonal adjustment reflects the plant's changing water use and nutrient demand patterns throughout the growing

Micronutrient Considerations

While macronutrients get most of the attention, micronutrient composition matters too. The (1) modified Steiner formulations included iron at 5 ppm, manganese at 1.6 ppm, boron at 0.865 ppm, zinc at 0.023 ppm, copper at 0.11 ppm, and molybdenum at 0.048 ppm. These concentrations remained constant across all treatments, suggesting that within reasonable limits, macronutrient balance has a more pronounced effect on yield and quality than micronutrient variation.

Making Practical Choices

So what should you actually do with this information? If you are growing strawberries hydroponically and want to maximize both yield and quality, consider starting with a solution containing approximately 160 to 170 ppm nitrogen, 55 to 60 ppm phosphorus, and 400 to 500 ppm potassium. Maintain the K:Ca ratio near 1-1.4:1 and the K:Mg ratio near 4:1. This matches some of my previous publications on the K:Ca ratio.

Remember that these recommendations assume you are maintaining appropriate pH (around 5.5 to 6.0) and EC levels suitable for your growing conditions. The (2) research demonstrated that excessive nutrients actually decreased both yield and quality, so more is definitely not better. You will need to adjust based on your specific cultivar, climate, and growing system, but these ranges provide a solid starting point backed by peer reviewed research.

The key takeaway is that strawberry nutrition in hydroponics requires a delicate balance. While nitrogen drives yield, potassium enhances quality, and the interaction between these two nutrients determines your ultimate success. Monitor your plants carefully, conduct tissue analysis when possible, and

do not be afraid to adjust your formulation based on what the plants are telling you.

Comparing Nutrient Solutions for Hydroponic Tomatoes

When growing tomatoes hydroponically, one of the most critical decisions you'll make is choosing the right nutrient solution. The composition of your nutrient solution can dramatically affect both the quantity and quality of your harvest. In this post, I'll examine different nutrient formulations that have been tested in scientific studies and discuss how they impact tomato production in soilless systems.



Picture of a soilless tomato greenhouse

Understanding Nutrient Solution Basics

Before diving into specific formulations, it's important to understand that tomato plants have changing nutritional needs throughout their growth cycle. Research has shown that early in the season, excessive nitrogen can cause plants to become too vegetative, resulting in bullish growth that produces misshapen fruits and increases susceptibility to disease (1). High potassium levels can also create problems by interfering with calcium and magnesium absorption, leading to blossom end rot.

Most successful nutrient programs divide the growing season into distinct stages. The seedling stage requires lower concentrations of nutrients, particularly nitrogen, while mature fruiting plants need substantially higher levels of most nutrients to support both vegetative growth and fruit development (2).

Comparing Two Common Formulations

Research has established several effective nutrient formulations for hydroponic tomatoes. I'll compare two well documented approaches that represent different philosophies in nutrient management.

Nutrient	Arizona Formula (Seedling)	Arizona Formula (Fruiting)	Florida Formula (Early)	Florida Formula (Late)
Nitrogen (N)	113 ppm	144 ppm	60 to 70 ppm	150 to 200 ppm
Phosphorus (P)	62 ppm	62 ppm	39 ppm	39 ppm

Potassium (K)	199 ppm	199 ppm	200 ppm	300 to 400 ppm
Calcium (Ca)	122 ppm	165 ppm	150 to 200 ppm	150 to 200 ppm
Magnesium (Mg)	50 ppm	50 ppm	48 ppm	48 ppm

The Arizona formulation (2) maintains relatively consistent macronutrient levels between growth stages, with only modest increases in nitrogen and calcium as plants mature. In contrast, the Florida approach (1) uses much lower nitrogen during early growth to prevent bullishness, then dramatically increases both nitrogen and potassium during fruit production.

Micronutrient Requirements

While macronutrients often receive the most attention, micronutrients are equally essential for healthy tomato production. These elements remain fairly constant throughout the growing cycle (2). Standard micronutrient concentrations for hydroponically grown tomatoes include iron at 2.5 ppm, manganese at 0.62 ppm, boron at 0.44 ppm, zinc at 0.09 ppm, copper at 0.05 ppm, and molybdenum at 0.06 ppm.

Micronutrient	Concentration (ppm)
Iron (Fe)	2.5
Manganese (Mn)	0.62
Boron (B)	0.44
Zinc (Zn)	0.09
Copper (Cu)	0.05
Molybdenum (Mo)	0.06

The Impact of Nitrogen Supply on

Quality

Research on nitrogen management has revealed some surprising findings. A study examining nitrogen supply at different growth stages found that increasing nitrogen from 140 to 225ppm during the vegetative stage increased protein, vitamin C, and sugar content in fruits (3). However, the effect on lycopene and beta-carotene depended heavily on the potassium supply during the reproductive stage.

Other research examining lower nitrogen levels has shown that minimal nitrogen supply can actually enhance lycopene content in tomato fruits, particularly when coupled with sufficient water supply (4). Studies in hydroponic culture have demonstrated that either the lowest or medium levels of nitrogen application produced the best lycopene content, suggesting that optimal nitrogen levels for antioxidant production may be lower than those for maximum yield.

Potassium's Role in Fruit Quality

Potassium plays a fundamental role in determining tomato fruit quality. Research has demonstrated that increasing potassium supply during the reproductive stage significantly enhances sugar concentration, vitamin C content, protein levels, lycopene, and beta-carotene in tomato fruits (3). The effect is particularly pronounced when potassium levels increase from 200 to 500ppm.

Another comprehensive study found that high proportions of potassium in the nutrient solution increased quality attributes including fruit dry matter, total soluble solids content, and lycopene content (5). However, these same researchers found that high proportions of calcium improved tomato fruit yield and reduced the incidence of blossom end rot, highlighting the importance of balancing these two nutrients.

Electrical Conductivity Management

One of the most innovative approaches to nutrient management involves carefully controlling the electrical conductivity (EC) of the nutrient solution. A study in closed NFT (Nutrient Film Technique) systems examined three different EC replacement set points: 5, 7.5, and 10 mS/cm (6). Remarkably, the highest EC replacement set point produced yields equivalent to lower EC treatments while significantly improving fruit quality.

The higher EC replacement threshold resulted in better dry matter content and total soluble solids in berries. Additionally, it demonstrated superior environmental sustainability by reducing total nutrients discharged into the environment by 37% compared to the medium EC treatment and 59% compared to the low EC treatment (6). This approach challenges conventional thinking about salinity stress in tomato production.

Calcium Management and Blossom End Rot

Calcium nutrition presents one of the most common challenges in hydroponic tomato production. Blossom end rot, characterized by dark lesions on the blossom end of fruits, results from calcium deficiency in developing fruits. However, this deficiency often occurs even when calcium levels in the nutrient solution appear adequate (1).

The problem frequently stems from antagonism between nutrients. Excessive potassium in the nutrient solution can interfere with calcium uptake by plant roots. This is particularly problematic early in the season when using premixed fertilizers that contain high potassium levels. Growers working with water containing less than 50 ppm calcium need to

be especially cautious about potassium concentrations.

To minimize blossom end rot, it's critical to maintain calcium levels between 150 and 200 ppm while keeping early season potassium levels moderate. Some growers supplement calcium nitrate with calcium chloride to increase calcium availability without adding more nitrogen. Each pound of calcium chloride (36% Ca) in 30 gallons of stock solution increases calcium concentration by approximately 14 ppm in the final nutrient solution when injected at a 1% rate (1).

Effects on Yield and Quality Parameters

The differences between nutrient formulations can significantly impact both yield and fruit quality. Research consistently shows that inadequate nitrogen during fruiting stages produces lower yields, though the fruits may have better sugar content and flavor. Conversely, excessive nitrogen can produce abundant foliage at the expense of fruit production (4).

Potassium levels have a pronounced effect on fruit quality parameters. Adequate potassium improves fruit firmness, color development, and sugar content (3). However, excessive potassium can lead to calcium and magnesium deficiencies that compromise both yield and quality.

The timing of nutrient adjustments also matters significantly. Studies have shown that gradually increasing nutrient concentrations as plants transition from vegetative to reproductive growth produces better results than sudden changes in formulation. Plants that experience consistent, appropriate nutrition throughout their lifecycle typically show improved yields and more uniform fruit quality (6).

Practical Considerations

When implementing a nutrient program, several practical factors deserve consideration. Water quality plays a fundamental role in determining how much of each nutrient to add. Wells in many regions naturally contain significant calcium and magnesium, sometimes providing 40 to 60 ppm calcium (1). These naturally occurring nutrients should be factored into your formulation calculations.

The pH of your nutrient solution also affects nutrient availability. Research has established that maintaining pH between 5.5 and 6.0 ensures optimal nutrient uptake (2). Water with high alkalinity requires acidification, which can be accomplished using phosphoric acid or sulfuric acid depending on your phosphorus requirements.

The type of hydroponic system you're using may also influence your nutrient concentrations. Systems requiring fewer daily irrigation cycles may need higher nutrient concentrations to ensure plants receive adequate nutrition. The general principle is that nutrient concentrations should be higher in systems with less frequent fertigation compared to those with continuous or very frequent feeding (1).

Advanced Management: The Transpiration-Biomass Ratio

One of the most sophisticated approaches to nutrient management involves calculating a recovery solution based on the transpiration-biomass ratio (6). This method recognizes that the relationship between water use and dry matter production changes throughout the growing cycle.

Research has shown that the transpiration-biomass ratio is high early in the crop cycle (approximately 300 liters per kilogram of dry weight), decreases during mid-season to a

relatively stable phase, and then increases again late in the season (up to 400 liters per kilogram). This pattern suggests that nutrient concentrations should be adjusted accordingly: lower concentrations in the first and last phases, and higher concentrations during the middle phase when biomass accumulation is most rapid.

Conclusion

Successful hydroponic tomato production requires careful attention to nutrient solution composition. While several proven formulations exist, the research clearly shows that no single approach works best for all situations. The Florida formulation with its conservative early nitrogen levels may be ideal for preventing bullishness in greenhouse production, while higher EC strategies can improve fruit quality in closed systems.

Key takeaways from the scientific literature include: maintain nitrogen between 60 and 70 ppm early in the season to prevent excessive vegetative growth, increase potassium substantially during fruiting to enhance quality parameters, keep calcium between 150 and 200 ppm throughout the season while monitoring potassium levels to prevent antagonism, and consider that higher EC values (up to even 10 mS/cm) may be feasible limits for nutrient solution replacement in recirculating systems.

Starting with a well researched base formulation and making careful adjustments based on plant response, tissue analysis, and your specific growing conditions provides the most reliable path to optimizing both yield and quality in your hydroponic tomato crop. The scientific evidence demonstrates that nutrient management is not a one-size-fits-all proposition, but rather a dynamic process that should respond to both plant developmental stage and environmental conditions.

pH vs Nutrient Availability: Rethinking the Classic Charts

If you've been around hydroponics long enough, you've probably seen the ubiquitous "pH vs nutrient availability" chart. It usually looks like a series of colored bars, each showing how available a nutrient supposedly is across a pH range. The bars are wide for some nutrients at certain pH values, narrow for others, and the chart often comes with a moral: keep your solution pH between 5.5 and 6.5.

I discussed some of these issues in a <u>previous post</u>, but it's worth revisiting them here with a clearer chart. The problem is that most of these charts trace back to soil agronomy research from the 1930s and 1940s. They're not based on solution chemistry relevant to hydroponics. They conflate microbial activity, lime chemistry, and plant physiology with solubility. And, in some cases, they are flat out misleading.

Let me talk about why the traditional chart is wrong, what modern chemistry tells us, and how a more honest representation looks.

Where the Old Charts Went Wrong

The historical diagrams were designed for soils, not hydroponic solutions. For example:

• Nitrate (NO₃⁻): In many charts, nitrate availability appears to fall off at low pH. In reality, nitrate is completely soluble across any reasonable pH range. The


- "loss" in those charts comes from soil microbial nitrification shutting down under acidic conditions, not relevant when you're directly dosing nitrate salts in solution.
- Calcium (Ca) and Magnesium (Mg): Old charts show Ca and Mg as always available at high pH. But that ignores precipitation with phosphate or carbonate, which can start as low as pH 6.2 for Ca. The old charts show high Ca and Mg availability at high pH because the high pH in soils was usually achieved by the addition of dolomite or lime, which greatly increased Ca and Mg concentrations in soil, this is not the case in a soilless setup.
- Micronutrients (Fe, Mn, Zn, Cu): These are shown as less available above neutral pH, which is true for unchelated forms (they hydrolyze and precipitate quickly). But in hydroponics, I typically use chelates, and their stability extends availability well above pH 7.
- Phosphorus (P): Charts often suggest a broad plateau around pH 6 to 7. In truth, phosphate solubility is sharply influenced by calcium concentration and carbonate alkalinity. The idea of a universal "wide bar" is misleading.

These errors matter. They lead growers to overemphasize the magic 5.5 to 6.5 range without appreciating that different nutrients behave differently, and that chelation or precipitation risks can change the picture entirely.

Building a Better Chart

To improve on the old diagrams, I constructed a new heatmap. Instead of arbitrary bar widths, each nutrient's relative availability (scaled from $\theta = low$ to l = high) is modeled

based on actual solubility, speciation, and chelation chemistry. The chart covers pH 4.0 to 8.5.



Updated chart I created for nutrient availability in soilless systems based on chemical and plant physiology principles

This chart is not an absolute quantitative prediction (real world systems have variations depending on concentration, alkalinity, chelate type, etc.). But it captures the directional chemistry more honestly. For nutrients that are effectively pH independent (like nitrate), the line is flat. For those that crash with pH (like unchelated iron), the line drops. And for Ca and Mg, I've introduced tapering to reflect phosphate precipitation behavior.

Nutrient by Nutrient Ranges

Here's a summary table describing the approximate pH behavior, the range of best availability, and the underlying reason:

Nutrient	Broad Availability Range	Notes / Reason
NO 3 ⁻ - N	4.0 to 8.5	Soluble across all relevant pH; uptake independent of pH in hydroponic solution. Old charts confused microbial nitrification with solubility.
NH 4 + - N	Best <6.5; declines >7.0	At higher pH, conversion to unionized NH3 increases, which is less available and potentially toxic.
Phosphorus (P)	Peak 5.5 to 6.5; drops <5.2 and >7.0	Solubility falls at high pH due to Ca+P precipitation (starting ~6.2); also limited at low pH by fixation and speciation.
Potassium (K)	4.0 to 8.5	Monovalent cation, highly soluble, minimal precipitation issues (sometimes K containing silicates at higher pH values)
Calcium (Ca)	Stable <6.0; declining >6.2	Precipitates with phosphate and carbonate as pH rises; availability falls gradually above ~6.2.
Magnesium (Mg)	Stable <6.5; mild decline >7.0	Mg+P precipitation is less aggressive than Ca+P; solubility loss is slower but still possible at higher pH.

Sulfate (SO4 ²⁻)	Broad 4.5 to 8.0	Generally soluble. At very low pH, some soils can adsorb sulfate due to protonated variable charge surfaces, reducing availability. At very high pH, reduced root uptake efficiency and competition with other anions can occur; in concentrated Ca ²⁺ + SO ₄ ²⁻ systems gypsum may precipitate by saturation.
Iron (Fe, unchelated)	Max <5.5; falls sharply >6.0	Fe ³⁺ hydrolyzes and precipitates as hydroxides and oxides above ~pH 6; nearly unavailable by pH 7.
Manganese (Mn, unchelated)	Best <6.0; declining >6.3	Mn ²⁺ oxidizes and precipitates above neutral pH.
Zinc (Zn, unchelated)	Best <6.0; low >7.0	Zn ²⁺ solubility decreases with increasing pH; precipitates as hydroxide/carbonate.
Copper (Cu, unchelated)	Best <6.0; poor >7.0	Cu ²⁺ strongly hydrolyzes, falls out of solution quickly with rising pH.
Boron (B)	Best 5.5 to 6.8	Boric acid is readily available in this range; at higher pH, more borate forms, reducing uptake.
Molybdenum (Mo)	Improves >6.0	Molybdate solubility increases with pH; plants often deficient in acidic conditions, more available at neutral/alkaline pH.

The Ca vs Mg Difference

A key improvement over older charts is distinguishing calcium from magnesium. While both can precipitate with phosphate, their behaviors differ:

- Ca+P precipitation is strong and begins around pH 6.2, especially in solutions with 1 to 3 mM phosphate. Brushite, dicalcium phosphate, and hydroxyapatite phases progressively reduce solubility.
- Mg+P precipitation is slower and less pronounced. Mg²⁺ is more strongly hydrated and less eager to form insoluble phosphates. It tends to stay soluble longer, only declining gently above pH 7.

Chelation: The Missing Dimension

My chart above shows unchelated forms. In real hydroponics, Fe, Mn, Zn, and Cu are almost always chelated. Depending on the chelate (EDTA, DTPA, EDDHA, HBED), stability can be maintained up to pH 7.5 to 9. This dramatically extends availability, particularly for Fe. A separate chart is needed to show chelated behavior.

Why This Matters

So why obsess about getting this chart right?

Because oversimplified charts lead to oversimplified thinking. If you believe nitrate solubility collapses below pH 6, you

might panic when your reservoir drifts to 5.2, even though NO_3^- is unaffected. If you believe Ca is "always available," you might miss that phosphate precipitation is happening in your tank right now at pH 6.3. And if you don't distinguish between chelated and unchelated micronutrients, you'll misdiagnose deficiencies.

A better chart isn't just about scientific pedantry. It's about helping growers make better decisions: when to acidify, when to buffer, when to choose a stronger chelate, and when to worry (or not worry) about a drifting pH.

Final Thoughts

The classic nutrient pH charts had their place in teaching basic agronomy 80 years ago. But hydroponics deserves more precision. Nutrients don't all behave the same way. Some are flat across the entire range $(NO_3^-,\ K)$. Some rise or fall gradually (B, Mo, Mg). Others are brutally sensitive (Fe without chelates). And precipitation interactions mean that Ca and phosphate availability are tied together, not independent.

This new heatmap and the accompanying table aren't the last word, they're a more honest starting point. The real message is: understand the chemistry, not just the cartoon.

Can you manage downy mildew in hydroponic basil with

organic foliar sprays?

mildew. by the Basil downv caused obligate oomycete Peronospora belbahrii, has become one of the most serious diseases affecting hydroponic and greenhouse basil production globally. The pathogen, first documented in Europe in 2001 and later detected in the United States in 2007, requires high relative humidity (at least 85%) or wet leaves to infect plants (1). Temperature preferences favor moderate conditions around 20°C rather than higher temperatures, which explains why the disease thrives in controlled environment systems where leaf wetness and humidity are difficult to manage (1).



Downy mildew in basil shows characteristic black marks on the underside of leaves

Understanding the infection process is critical for designing effective spray programs. Under conditions of continuous free moisture, sporangia germinate within 3 to 5 days by producing germ tubes that penetrate basil leaves directly through the epidermis, typically without entering through stomata (2). Seven days after initial infection, sporangiophores bearing new sporangia emerge through stomata on both the upper and lower leaf surfaces, creating secondary inoculum that spreads rapidly throughout greenhouse facilities (2). This relatively

short cycle from infection to sporulation means that preventive measures must start before visible symptoms appear.

Multiple field trials evaluating organic fungicides have delivered sobering results for growers seeking alternatives to conventional chemistry. A comprehensive study testing products approved for organic production, including copper octanoate, hydrogen dioxide, sesame oil, neem oil, thyme oil, citric acid, Bacillus species, and Streptomyces lydicus, found that none were effective at controlling downy mildew when applied to susceptible basil cultivars (3). Applications were made weekly starting before symptom development, and efficacy was assessed based on incidence of symptomatic leaves rather than severity, reflecting the zero tolerance for disease on fresh market herbs (3). A summary of the tested fungicides and their effectiveness is shown on the following table.

Product (Active Ingredient)	Mode of Action	Effectiveness
Cueva (Copper octanoate)	Contact fungicide, disrupts enzyme function	Ineffective
OxiDate (Hydrogen dioxide)	Oxidizing agent, contact action	Ineffective
Organocide (Sesame oil)	Physical barrier, suffocation	Ineffective
Trilogy (Neem oil)	Physical barrier, azadirachtin content	Ineffective
Forticept EP #1 (Thyme oil)	Essential oil, contact action	Ineffective
Procidic (Citric acid)	pH modulation, contact action	Ineffective
Actinovate (Streptomyces lydicus)	Biocontrol, competitive colonization	Ineffective

Product (Active Ingredient)	Mode of Action	Effectiveness
Companion (Bacillus subtilis)	Biocontrol, induced resistance	Ineffective
Double Nickel (B. amyloliquefaciens)	Biocontrol, antibiosis	Ineffective
Regalia (Reynoutria sachalinensis)	Plant defense activator	Ineffective

The limited efficacy of organic fungicides appears related to the aggressive nature of the pathogen and the difficulty of achieving thorough foliar coverage in dense basil canopies. Even when combined with resistance inducers or natural products, organic treatments failed to provide commercially acceptable levels of disease suppression (5).

Environmental management offers more promise than chemical sprays alone. Light suppresses sporulation of *P. belbahrii*, with continuous light or supplemental lighting during nighttime hours substantially reducing spore production (6). Growers can exploit this by maintaining photoperiods longer than 13 hours or by using low-intensity supplemental lighting during dark periods. Reducing leaf wetness duration is equally important because the pathogen requires at least 24 hours of continuous moisture for infection and dense sporulation (7). In hydroponic systems, switching from overhead misting to subcanopy irrigation and increasing air movement with horizontal airflow fans can dramatically reduce infection pressure (8).

Temperature manipulation provides another non-chemical tool. Passive heat treatment using transparent plastic covers to raise greenhouse temperatures during sunny periods suppressed downy mildew development without damaging basil plants (9). Temperatures above 30°C inhibit sporangiophore formation and sporangial germination, though plants must be acclimated gradually to avoid heat stress. This approach works best in greenhouse operations with sufficient ventilation control and

may be less practical in open hydroponic facilities.

Varietal resistance remains the most effective long-term strategy for hydroponic basil growers. Breeding efforts have identified resistance sources in wild basil species *Ocimum americanum*, and these traits have been successfully transferred into sweet basil backgrounds (10). Commercial varieties with improved resistance are now available, though complete immunity has not been achieved. Growers should prioritize these resistant cultivars and combine them with environmental controls rather than relying on organic fungicide sprays.

Cropping system modifications can reduce disease pressure in organic systems. Research on open field organic production found that sparse sowing density combined with resistant varieties provided better control than chemical treatments alone (11). In hydroponics, maintaining wider plant spacing, particularly in NFT or DWC systems where humidity tends to be higher, allows better air circulation and faster leaf drying after irrigation events.

The reality for hydroponic basil producers is that organic foliar sprays, when used alone, will not provide adequate downy mildew control on susceptible varieties. The pathogen's rapid lifecycle, preference for humid greenhouse conditions, and resistance to contact fungicides makes chemical intervention largely ineffective without supporting measures. Successful organic management requires integrating resistant varieties, environmental manipulation (particularly light, humidity, and leaf wetness control), appropriate plant spacing, and vigilant monitoring for early disease detection. Growers who continue relying primarily on organic sprays should expect continued losses, while those who adopt integrated approaches combining genetics and environment will achieve better results.