

Bio-stimulants: Which Pure Compounds Have Reproducible Effects

If you have been exploring ways to improve crop performance in your hydroponic system, you have likely encountered the term “bio-stimulants.” The market is flooded with products making bold claims, but separating marketing hype from reproducible science can be challenging. In this post I am going to focus exclusively on **pure chemical compounds** that have demonstrated consistent effects in peer-reviewed research. I am deliberately excluding mixtures, proprietary blends, polymeric substances, and commercial formulations to help you understand which individual substances actually work.

After reviewing the scientific literature extensively, I have identified several categories of pure bio-stimulants with strong evidence from multiple independent studies: specific amino acids, silicon compounds, plant hormones, melatonin, and thiamine. Each compound discussed below has at least five peer-reviewed studies demonstrating consistent positive effects in controlled greenhouse or hydroponic systems.



Taken from [this article](#), it shows the effect of some bio-stimulants, including melatonin, on calendula officinalis (one of my favorite plants). A layout of the experiment. Salinity levels (S), S0 = Tap water, S1 = 42.8 mM, S2 = 85.6 mM, S3 = 128.3 mM, Melatonin (M), M0 = 0 μM, M1 = 50 μM, M2 = 100 μM, Bacterial inoculation (B), B0 = non-inoculation, B1 = inoculation

What about humic and fulvic acids?

Before diving into the compounds that made the cut, I want to address a common question. Humic and fulvic acids are popular in hydroponics, but they do not qualify as pure substances. According to the International Humic Substances Society, these are “complex and heterogeneous mixtures of polydispersed

materials” containing thousands of distinct organic compounds (1). Modern analytical chemistry has identified **5,000 to 7,000 unique molecules** in typical humic extracts. While they can be effective bio-stimulants, they fall outside the scope of this article because their variable composition makes reproducibility difficult to guarantee across different sources.

Amino acids with extensive research support

Two amino acid compounds stand out for having robust evidence across multiple independent studies: **glycine betaine** and **L-proline**.

Glycine betaine functions as an osmoprotectant, stabilizing protein structure and protecting photosystem II under stress conditions (2). Commercial greenhouse hydroponic lettuce production in Finland demonstrated reduced nitrate accumulation while maintaining yield (3). Hydroponic trials in chickpea showed significant improvements in chromium stress tolerance at 11715 ppm (4). Field applications at 700 ppm improved lettuce performance under water stress (5). Pot studies with maize demonstrated enhanced growth and chlorophyll content under drought at concentrations of 3650 to 3840 ppm (6). Hydroponic maize trials with 11.7 ppm showed improved salt tolerance through Na⁺ homeostasis regulation (7). Field trials in winter wheat at 5858 ppm demonstrated improved water use efficiency under limited irrigation (8).

L-proline operates through similar osmoprotective mechanisms while also acting as a reactive oxygen species scavenger. Greenhouse hydroponic studies in maize showed significant drought tolerance improvements at 576 to 1151 ppm application rates (9). Field trials conducted in Egypt during 2017-2018 demonstrated that foliar proline at 230 to 461 ppm significantly improved maize yield under drought stress with

both surface and drip irrigation systems ([10](#)). Greenhouse tomato trials showed that 100 ppm proline application alleviated heat stress damage and increased fruit yield per plant ([11](#)). Tomato seedling studies demonstrated that 1151 ppm foliar proline provided protection against chilling stress through enhanced antioxidant enzyme activities ([12](#)). Hydroponic NFT tomato trials with 1151 ppm foliar proline application alleviated salinity stress effects on cell ultrastructure and photosynthesis ([13](#)). Multiple greenhouse studies confirmed proline improved stress tolerance across various crops at concentrations between 576 to 2878 ppm ([14](#)).

Silicon: the most extensively validated bio-stimulant

Potassium silicate (K_2SiO_3) is the most practical option for nutrient solution supplementation. At hydroponic pH levels, it hydrolyzes into monosilicic acid and potassium ions. Plants absorb the monosilicic acid through specialized aquaporin-type channels and deposit it as amorphous silica in cell walls ([15](#)). This creates physical barriers against pathogens while improving structural integrity.

An important point to understand about silicon sources: at the pH where plants are fed in hydroponics, acid-stabilized silicon products and potassium silicate sources generate the exact same monosilicic acid. Stabilized monosilicic acid products are not more plant available than potassium silicate. The advantage of stabilized products is that they remain stable longer in recirculating systems and do not require pH adjustment, while potassium silicate polymerizes relatively quickly at typical hydroponic pH values.

Multiple greenhouse trials demonstrated pronounced resistance to powdery mildew in cucumber at 477 ppm Si ([16](#)). Melon greenhouse studies showed 65 to 73 percent reduction in powdery mildew disease progress with root application ([17](#)).

Hydroponic barley trials at various concentrations confirmed growth improvements ([18](#)). Greenhouse cucumber studies demonstrated that silicon addition to nutrient solutions significantly reduced powdery mildew severity ([19](#)). Recent lettuce research showed silicon extended shelf life by 40 to 80 percent ([20](#)). Zucchini greenhouse trials confirmed silicon effectiveness against powdery mildew when applied both foliar and through roots ([21](#)).

Melatonin: an emerging bio-stimulant with strong evidence

Melatonin has emerged as a promising bio-stimulant with extensive research support across multiple crops. This compound functions as both an antioxidant and growth regulator.

Hydroponic tomato trials demonstrated that 11.6 to 46.5 ppm melatonin improved growth and photosynthetic characteristics under saline-alkali stress ([22](#)). Greenhouse cucumber studies at 23.2 ppm showed enhanced nitrogen metabolism and growth ([23](#)). Tomato fruit quality studies confirmed that 23.2 ppm melatonin promoted accumulation of sugars, amino acids, and secondary metabolites ([24](#)). Hydroponic wheat trials with 23.2 ppm enhanced drought tolerance through jasmonic acid and lignin bio-synthesis pathways ([25](#)). Cucumber seed priming with melatonin improved antioxidant defense and germination under chilling stress ([26](#)). Greenhouse tomato trials demonstrated that 116 ppm melatonin improved salt tolerance when applied as foliar spray ([27](#)). Multiple studies confirmed melatonin at 11.6 to 116 ppm enhanced photosynthesis, antioxidant systems, and stress tolerance across various crops ([28](#)).

Thiamine (Vitamin B1): disease

resistance activator

Thiamine has a unique position among bio-stimulants due to its role in activating systemic acquired resistance in plants rather than direct nutritional effects.

Greenhouse studies demonstrated that foliar application of 5772 ppm thiamine induced systemic acquired resistance in rice, Arabidopsis, tobacco, and cucumber against fungal, bacterial, and viral infections (35). Wheat pot trials showed that 100 ppm thiamine improved growth, chlorophyll content, and yield under water stress (36). Research confirmed thiamine functions as an activator of plant disease resistance through salicylic acid and calcium-dependent signaling pathways (35). Greenhouse trials on multiple crops demonstrated that thiamine treatment at 50 to 100 ppm protects plants against biotic and abiotic stresses (37). Studies showed thiamine enhanced stress tolerance by improving thiamine bio-synthesis pathway regulation under osmotic and salt stress (37). Research on various plant species confirmed thiamine involvement in primary metabolism and stress response mechanisms (38). Soybean trials demonstrated that 50 to 100 ppm thiamine favors plant development and grain yield as a bio-stimulant (39).

Important note: Thiamine does NOT stimulate root growth or reduce transplant shock in whole plants despite common marketing claims. Its beneficial effects are limited to disease resistance and metabolic enhancement.

Plant hormones with consistent small-scale validation

Gibberellic acid (GA3) has extensive greenhouse and laboratory validation across multiple crops. Hydroponic lettuce and rocket floating system trials established tested concentrations around 0.35 ppm for enhanced growth and yield

(29). Hydroponic lettuce studies with 20 to 100 ppm GA3 showed improved morphological characteristics and yield (30). Greenhouse tomato seed treatment studies demonstrated that 300 to 900 ppm GA3 increased germination percentage and seedling vigor (31). Greenhouse trials on yellow cherry tomatoes showed that 25 to 75 ppm GA3 foliar applications increased stem diameter, branch number, and fruit biomass by up to 93.8% (32). Hydroponic cucumber studies confirmed that 1.7 ppm GA3 reversed growth inhibition caused by low root-zone temperatures (33). Greenhouse tomato seedling trials demonstrated that GA3 treatment improved growth and reduced heavy metal accumulation under stress conditions (34). The compound decreased nitrate accumulation in leafy vegetables while increasing dry weight. Concentrations around 0.35 ppm are widely used in research settings for various crops, though higher concentrations cause excessive elongation that reduces marketability.

Salicylic acid shows consistent benefits across greenhouse trials. Hydroponic cucumber studies demonstrated yield improvements at 69 ppm (40). Greenhouse tomato trials showed positive effects on plant growth and yield at 69 ppm applications (41). Greenhouse tomato trials with 250 ppm salicylic acid enhanced drought tolerance through improved antioxidant enzyme activity (42). Field tomato studies demonstrated 40 to 45 percent yield increases at 138 to 207 ppm under water stress (43). Greenhouse cucumber trials confirmed improved phenolic compounds and yield at 10.4 to 69 ppm (44). Hydroponic maize studies showed protection against chilling injury at 69 ppm (45).

Suggested test application rates and practical suggestions

Based on the evidence reviewed, here are some suggestions if you want to try pure compound bio-stimulants. As always, make

sure to try on a small number of plants before making large scale applications:

For silicon supplementation, potassium silicate at 20 ppm Si (approximately 40 ppm SiO_2) offers excellent disease resistance and yield benefits. Add it to your nutrient solution at each reservoir change and adjust pH accordingly. Remember that low cost potassium silicates can provide readily available monosilicic acid when used properly. For more details on silicon use in hydroponics, see [this previous article](#).

For stress tolerance, glycine betaine at 700 ppm in nutrient solution or L-proline at 575 ppm as foliar application can significantly improve crop performance under salt or drought conditions. For comprehensive guidance on glycine betaine applications, see [this previous article](#).

For melatonin applications, use 25 ppm as foliar spray or in nutrient solution. This concentration has shown consistent benefits across multiple crops for stress tolerance and growth enhancement.

For disease resistance, thiamine at 100 ppm as foliar spray activates systemic acquired resistance. This is particularly useful for preventive disease management rather than direct growth promotion. For detailed information on thiamine applications, see [this previous article](#).

For specialized applications, gibberellic acid at 0.35 ppm or salicylic acid at 30 ppm offer targeted benefits, though these require more careful application timing and concentration control. For more information on salicylic acid use, see [this previous article](#).

Summary table: Pure compounds with

reproducible effects

Compound	Number of Studies	Tested Concentration	Primary Benefits
Glycine Betaine	7 studies (2 , 3 , 4 , 5 , 6 , 7 , 8)	12–5900 ppm	Osmoprotection, salt tolerance, reduced nitrate
L-Proline	6 studies (9 , 10 , 11 , 12 , 13 , 14)	230–2900 ppm (foliar)	ROS scavenging, drought tolerance, salt stress
Potassium Silicate	7 studies (15 , 16 , 17 , 18 , 19 , 20 , 21)	14–42 ppm Si	Disease resistance, shelf life, structural integrity
Melatonin	7 studies (22 , 23 , 24 , 25 , 26 , 27 , 28)	11–116 ppm	Antioxidant activity, stress tolerance, growth regulation
Gibberellic Acid	6 studies (29 , 30 , 31 , 32 , 33 , 34)	0.35–1.7 ppm	Fruit development, reduced nitrate, cell elongation
Thiamine (Vitamin B1)	5 studies (35 , 36 , 37 , 38 , 39)	50–100 ppm (foliar)	Disease resistance activation, stress metabolism
Salicylic Acid	6 studies (40 , 41 , 42 , 43 , 44 , 45)	70–250 ppm	Stress tolerance, yield enhancement, disease resistance

The key advantage of using pure compounds rather than commercial blends is reproducibility. When you know exactly what you are applying and at what concentration, you can systematically optimize your system and troubleshoot problems effectively. Each of these compounds has been validated across multiple independent studies, giving you confidence that results can be consistent across different growing conditions.

However, keep in mind that crop conditions can be very variable and, while these bio-stimulants have been validated across various scenarios, effects can vary depending on the particular circumstances of each crop.

Have you tried any of these pure compound bio-stimulants in your hydroponic system? What were your results? Let us know in the comments below!

How to prevent problems with powdery mildew in hydroponic crops

One of the worst problems you can get in a hydroponic crop is mildew. Year after year I see growers lose significant amounts of production due to this disease within a variety of different crops. Powdery mildew reduces yields, stunts plants and – if contracted early on – will possibly cause a complete loss of your crop. It is generally hard to control once it gets in and it will expand like wildfire through any commercial growing operation. Today we will be discussing how to actually prevent mildew from ever appearing – without using toxic fungicide applications – and why prevention can play a huge role in ensuring you never have to face this problem in

the first place.



Fungal spores are generally carried by the wind and by insects, making it very hard for a crop to avoid ever coming into contact with the pathogen. Wild plants or plants from other commercial crops close to you will most likely have the disease and millions of spores will get in the air and eventually reach your plants. It is only a matter of time till the powdery mildew reaches your crops – almost impossible to prevent – so you must make sure that your plants are strong enough to prevent the pathogen from taking hold.

There are two main factors that affect whether powdery mildew will infect your plants. The first is plant strength and the second is the environment. If one of these two is not at its best then your plants will fall prey to this fungal disease. Neither strong plants under bad environmental conditions nor weak plants under ideal environmental conditions will be safe from the disease. So what can we do to ensure our plants are healthy and our environmental conditions are safe?

One of the proven methods to make plants strong against fungi is silicon. Potassium silicate applications – as soil drenches or foliar sprays – have proven to increase disease resistance across several studies (see [here](#) and [here](#) for examples). But other innovative approaches using other forms of silicon – for example nanometer sized silica crystals – have also yielded good results. In [this](#) and [this](#) studies it was clearly shown that other forms of silicon – besides silicate – could also help in preventing fungal disease. This might be preferred in some cases as these forms of silicon can be far more stable and easier to store/apply compared with options like potassium silicate.





However silicate applications are no miracle. If your environmental conditions are not set properly the silicate applications will be useless. This is the reason why some growers say that silicate does nothing against disease, because an environment that's favorable for fungi can basically nullify the protective action of supplemental silicon. This was demonstrated by cucumber growers who had a lot of success with Si supplementation in Canada to prevent fungal diseases, but failed to reproduce this success in Florida. A [study about this](#) difference revealed that the higher temperatures in Florida negated a large part of the benefits from silicon supplementation. If you want silicon to work against disease better stay in the 20-25°C range.

Other microorganisms can also be of great help in preventing powdery mildew. If a leaf is already colonized by beneficial fungi or bacteria it will be much harder for a pathogen to get in. Several species of microorganisms have been studied in this regard. Fungi like *Tilletiopsis* have shown to prevent and control the disease (see [here](#)), other microbes have also been studied in conjunction with silicon (see [here](#) and [here](#)), showing beneficial effects. Fungus like *Trichoderma harzianum* and bacteria like *Bacillus subtilis* have also shown induction of systemic resistance against fungal diseases (see [here](#), [here](#) and [here](#)). The two images above were taken from [this study](#).

Friendly chemical solutions are also available for the prevention of powdery mildew. Plant derived extracts, for example neem seed oil at 1% has shown to be a good agent for powdery mildew prevention in okra (see [here](#)). Substances like salicylic acid have also shown to trigger resistance to powdery mildew in plants like peas (see [here](#)).



There are also additional alternatives dealing with the

environment that can make it difficult for fungi to colonize plants by attempting to make the environment more hostile for fungi. Spraying ozonated water has shown positive results in experiments with tomatoes (see [here](#)) as well as electrolyzed water in strawberries (see [here](#)). Keeping the environment conditions within a proper range is also important, [this paper](#) shows you how environmental conditions affect powdery mildew disease severity in sunflower but the general features are applicable to most higher plants. As you can see in the image above – taken from this paper – disease severity increases with relative humidity. In general you will want to keep your relative humidity below 70% to avoid making the environment extremely friendly for fungi.

In the end there are many things you can do to keep your plants free of foliar fungal disease like powdery mildew. Use lower temperatures, control your relative humidity, do silicate and salicylic acid applications and use beneficial microbes. If you follow these steps you will forget that powder mildew ever existed!