


# Triaccontanol Foliar Sprays in Soilless Culture: Formulation and Application

Triaccontanol is a naturally occurring long-chain fatty alcohol found in plant cuticle waxes that can act as a growth regulator at very low concentrations. Below I focus on peer-reviewed evidence for triaccontanol in hydroponic and soilless systems, with attention to preparation methods, yield effects, and quality outcomes in tomatoes, cucumbers, strawberries, and lettuce.



Above you can see a representative model of triaccontanol. Chemically triaccontanol is a long-chain fatty alcohol, very hard to dissolve in water and apply effectively to plants.

## Evidence for Yield and Quality Effects

**Hydroponic lettuce.** Foliar application of triaccontanol at  $10^{-7}$  M (approximately 0.043 mg/L) to 4-day-old hydroponically grown lettuce seedlings increased leaf fresh weight by 13-20% and root fresh weight by 13-24% within 6 days. [\(1\)](#) When applied at both 4 and 8 days after seeding, leaf area and mean relative growth rate increased by 12-37%. There was no additional benefit from repeating applications beyond two sprays in this short-cycle crop.

**Tomato in hydroponic systems.** Weekly foliar applications of 70  $\mu$ M triacontanol (approximately 21 mg/L) on tomatoes grown in hydroponic drip systems significantly increased flower number by 37-50% and total fruit number by 22-57%, resulting in a 28% higher total yield at harvest. [\(2\)](#) Individual fruit weight decreased by 16%, but the net effect on total productivity remained positive. The treatment advanced blooming without affecting plant height or internode number, demonstrating a specific effect on reproductive development.

**Cucumber under soilless conditions.** Foliar application of triacontanol at 0.8 mg/L on cucumber genotypes under salt stress improved photosynthesis, stomatal conductance, and water use efficiency. [\(3\)](#) The treatment enhanced antioxidant enzyme activities and maintained better membrane stability. Yield traits, including fruit number and average fruit weight, improved in response to triacontanol application. Salt-tolerant genotypes (Green long and Marketmore) showed greater responsiveness than sensitive genotypes.

**Strawberry.** Triacontanol has shown promise in improving drought tolerance in strawberry plants by enhancing growth, productivity, and physiological performance, though most work has been conducted in soil rather than true soilless systems. [\(4\)](#)

## **Formulation:                      Creating                      a Concentrated Stock Solution**

Triacontanol has extremely low water solubility (less than 1 mg/L at room temperature), which makes proper formulation critical. The most reliable approach combines an organic solvent with a surfactant to create a stable concentrate that can be diluted into spray solutions.

# Stock Solution Protocol

## Materials needed:

- Triacontanol powder (90%+ purity)
- Ethanol (95% or higher)
- Tween-20 or Tween-80 (polysorbate surfactant)
- Distilled or deionized water
- Glass or high-density polyethylene containers

## Preparation of 1000 mg/L (1000 ppm) stock:

1. Weigh 1000 mg of triacontanol powder using an analytical balance.
2. Dissolve the triacontanol in 100 mL of 95% ethanol in a glass beaker. Warm gently (35-40°C) while stirring with a magnetic stirrer for 15-20 minutes to ensure complete dissolution. Do not exceed 50°C.
3. Add 5 mL of Tween-20 to the ethanol solution and mix thoroughly for 5 minutes. This surfactant concentration (0.5% v/v in final volume) ensures proper emulsification and leaf surface wetting.
4. Transfer the ethanol-triacontanol-surfactant mixture to a 1000 mL volumetric flask.
5. Bring to final volume with distilled water while mixing continuously. The solution will appear slightly cloudy due to micelle formation, which is expected and desirable.
6. Store the stock solution in an amber glass bottle at room temperature. The stock is stable for 3-4 months when protected from light and heat.

**Alternative solvent systems:** Some studies have successfully used isopropanol or acetone as solvents. [\(5\)](#) However, ethanol provides the best combination of triacontanol solubility, plant safety, and ease of handling for growers.

# Working Solution Preparation

Dilute the 1000 mg/L stock to achieve target concentrations based on crop and growth stage:

**Lettuce:** Dilute 1:10,000 to 1:20,000 for final concentrations of 0.05-0.1 mg/L. For a 1-liter spray bottle, add 0.05-0.1 mL of stock solution.

**Tomato:** Dilute 1:50 for final concentration of 20 mg/L. For a 1-liter spray bottle, add 20 mL of stock solution.

**Cucumber:** Dilute 1:1250 for final concentration of 0.8 mg/L. For a 1-liter spray bottle, add 0.8 mL of stock solution.

Add an additional 0.1% v/v Tween-20 (1 mL per liter) to the final spray solution to ensure maximum leaf coverage and absorption. This additional surfactant enhances uptake without phytotoxicity when concentrations remain below 0.2%. [\(3\)](#)

## Application Timing and Frequency

**Seedling stage:** Apply once at 4-8 days after emergence for leafy greens in short-cycle production. A single early application is often sufficient for lettuce. [\(1\)](#)

**Vegetative and reproductive stages:** For fruiting crops like tomato and cucumber, apply weekly starting 4 weeks after transplant and continuing through flowering and early fruit set. Three to five applications total are typically used. [\(2\)](#)  
[\(3\)](#)

**Application method:** Apply using a hand sprayer or backpack sprayer with a cone nozzle, ensuring complete leaf coverage including undersides. Apply in early morning or late afternoon to maximize absorption and minimize evaporation. Spray until runoff just begins.

# Reported Effects Across Crops

Crop	Concentration	Application schedule	Yield effect	Quality effect	Reference
Lettuce (hydroponic)	0.043 mg/L	Once at day 4, optional repeat at day 8	Fresh weight +13-20%, leaf area +12-37%	Not assessed	<a href="#">(1)</a>
Tomato (hydroponic drip)	21 mg/L	Weekly from week 4 through fruit set	Total yield +28%, fruit number +22-57%	Minimal changes in soluble solids, lycopene, vitamin C	<a href="#">(2)</a>
Cucumber (soilless, salt stress)	0.8 mg/L	Three sprays: 72h after stress, at flowering, at fruit maturity	Improved fruit number and weight under stress	Maintained lower electrolyte leakage, higher chlorophyll	<a href="#">(3)</a>

## Mechanisms and Considerations

Triacantanol acts through a secondary messenger system involving 9-L(+)-adenosine, which triggers rapid ion influx ( $\text{Ca}^{2+}$ ,  $\text{K}^{+}$ ,  $\text{Mg}^{2+}$ ) and modulates gene expression related to photosynthesis, hormone balance, and stress responses. [\(2\)](#) The compound enhances photosynthetic rate, stomatal conductance, and nutrient uptake at very low doses.

Concentration matters. Response curves show classic hormesis: stimulation at low concentrations, no effect or inhibition at higher doses. The optimal range is crop-specific but generally falls between 0.05-20 mg/L for foliar applications. Lettuce seems to respond to much lower concentrations than tomatoes.

Environmental and genetic factors influence response magnitude. Tolerant cucumber genotypes showed larger yield improvements than sensitive ones. [\(3\)](#) Season, light intensity, and nutrient status affect outcomes.

Triacontanol enhances stress tolerance, particularly to salinity and drought, by improving antioxidant enzyme activity, maintaining membrane integrity, and regulating osmotic adjustment. [\(3\)](#) [\(4\)](#) This makes it especially valuable in recirculating hydroponic systems where EC can drift upward.

## Practical Guidelines

- Test on a small number of plants before scaling to full production.
- Keep application rates within published ranges. More is not better with triacontanol.
- Maintain consistent spray timing rather than irregular high-dose applications.
- Store stock solutions away from light and heat to preserve activity.
- Use analytical-grade triacontanol from reputable suppliers (minimum 90% purity).
- Combine with sound nutritional management; triacontanol is not a substitute for balanced feeding. Triacontanol is not a replacement for proper nutrition, irrigation, environmental conditions or media management.

Properly formulated and applied, triacontanol provides measurable improvements in productivity and stress tolerance across major soilless crops. The citations above offer detailed protocols and results for those wishing to implement this growth regulator in commercial or research settings.